

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^{Pf}LEUCOVORIN CALCIUM INJECTION USP

Leucovorin calcium

Solution, 10 mg/mL, intravenous and intramuscular

USP

Folic Acid Derivative

Pfizer Canada ULC
17300 Trans-Canada Highway
Kirkland, Québec
H9J 2M5

Date of Initial Authorization:
MAY 09, 1997

Date of Revision:
DEC 13, 2022

Submission Control Number: 267164

RECENT MAJOR LABEL CHANGES

7 WARNINGS AND PRECAUTIONS, Reproductive Health: Female and Male Potential; 7.1.1 Pregnant Women	12/2022
--	---------

Table of Contents

RECENT MAJOR LABEL CHANGES	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	5
4 DOSAGE AND ADMINISTRATION	5
4.2 Recommended Dose and Dosage Adjustment	5
4.3 Reconstitution	7
4.4 Administration	7
5 OVERDOSAGE	8
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	8
7 WARNINGS AND PRECAUTIONS	8
7.1 Special Populations	10
7.1.1 Pregnant Women	10
7.1.2 Breast-feeding	10
7.1.3 Pediatrics	10
7.1.4 Geriatrics	11
8 ADVERSE REACTIONS	11
8.1 Adverse Reaction Overview	11
8.2 Clinical Trial Adverse Reactions	11
8.5 Post-Market Adverse Reactions	12
9 DRUG INTERACTIONS	12
9.1 Serious Drug Interactions	12

9.4	Drug-Drug Interactions	12
9.5	Drug-Food Interactions	14
9.6	Drug-Herb Interactions	14
9.7	Drug-Laboratory Test Interactions.....	14
10	CLINICAL PHARMACOLOGY	14
10.1	Mechanism of Action	14
10.2	Pharmacodynamics.....	14
10.3	Pharmacokinetics.....	14
11	STORAGE, STABILITY AND DISPOSAL.....	15
12	SPECIAL HANDLING INSTRUCTIONS.....	15
PART II: SCIENTIFIC INFORMATION		16
13	PHARMACEUTICAL INFORMATION	16
14	CLINICAL TRIALS	16
15	MICROBIOLOGY	16
16	NON-CLINICAL TOXICOLOGY	16
PATIENT MEDICATION INFORMATION		18

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Leucovorin Calcium Injection USP (leucovorin calcium) is indicated:

- To diminish the toxicity and counteract the effects of overdosage of folic acid antagonists.
- To diminish the systemic toxicity of methotrexate after administration of methotrexate as a chemotherapeutic agent, as part of chemotherapeutic treatment programs in the management of several forms of cancer.
- To treat megaloblastic anemias due to folate deficiency, as in sprue and other nutritional deficiencies; and megaloblastic anemias of pregnancy and infancy (see 2 [CONTRAINDICATIONS](#)).
- For pre-treatment followed by fluorouracil to prolong survival in the palliative treatment of patients with advanced colorectal cancer.

1.1 Pediatrics

Pediatrics (< 18 years of age): No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use (see 7.1.3 [Pediatrics](#)).

1.2 Geriatrics

Geriatrics: Evidence from clinical studies and experience suggests that use in the geriatric population is associated with differences in safety or effectiveness (see 7 WARNINGS AND PRECAUTIONS, General and 7.1.4 [Geriatrics](#)).

2 CONTRAINDICATIONS

Leucovorin Calcium Injection USP is contraindicated for:

- Pernicious anemia therapy or other megaloblastic anemias secondary to a deficiency of vitamin B₁₂. Its use can lead to an apparent response of the hematopoietic system, but neurological damage may occur or progress if already present.
- Known hypersensitivity to this drug or to any ingredient in the formulation or component of the container (see 6 [DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#)).
- Intrathecal administration.

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings and Precautions

- Leucovorin Calcium Injection USP should only be given by intramuscular or intravenous injection and must not be administered intrathecally (see 2 [CONTRAINDICATIONS](#) and 4 [DOSAGE AND ADMINISTRATION](#)). Death has been reported when folinic acid has been administered intrathecally following intrathecal overdose of methotrexate.
- Leucovorin Calcium Injection USP should only be used with 5-fluorouracil or methotrexate under the direct supervision of a clinician experienced in the use of cancer chemotherapeutic agents.
- Patients receiving any combination therapy regimen involving leucovorin and fluorouracil should be carefully monitored for diarrhea and/or stomatitis/mucositis as these are the first indications that severe and potentially life-threatening toxicity could develop (see 7 [WARNINGS AND PRECAUTIONS](#) and 9 [DRUG INTERACTIONS](#)).
- Fatalities have occurred as a result of gastrointestinal toxicity (predominantly mucositis and diarrhea) (see 7 [WARNINGS AND PRECAUTIONS](#) and 8 [ADVERSE REACTIONS](#)).
- Fatalities have occurred as a result of myelosuppression (see 7 [WARNINGS AND PRECAUTIONS](#) and 8 [ADVERSE REACTIONS](#)).
- Cases of Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN), some fatal, have been reported in patients receiving leucovorin in combination therapy (see 8 [ADVERSE REACTIONS](#)).
- Leucovorin may diminish the effect of anti-epileptic substances such as phenobarbital, primidone and phenytoin. During leucovorin administration in epileptic patients treated with these substances, there is a risk to increase the frequency of seizures due to a decrease of plasma concentrations of anti-epileptic drugs (see 9 [DRUG INTERACTIONS](#)).

4 DOSAGE AND ADMINISTRATION

4.2 Recommended Dose and Dosage Adjustment

Due to calcium content of leucovorin solution, no more than 160 mg of leucovorin should be injected, per minute, intravenously.

Treatment of Overdosage of Folic Acid Antagonists:

In cases of overdosage of folic acid antagonists, prompt administration of leucovorin calcium is essential; if a period of more than four hours intervenes, the treatment may be ineffective.

The dose of leucovorin calcium should be equal to or greater than the suspected dose of folic acid antagonist.

Where large doses of methotrexate have been given, leucovorin may be administered by intravenous infusion in doses up to 75 mg within 12 hours, followed by 12 mg intramuscularly every six hours, for four doses. In less severe overdosage, 6 to 12 mg of leucovorin may be given intramuscularly every six hours, for four doses, until the serum methotrexate level is less than 10^{-8} M.

Use After Chemotherapy with Methotrexate:

The dosage and scheduling of doses of leucovorin varies, but it is normally given about 6 to 24 hours following methotrexate administration, in amounts equal to the weight of methotrexate given.

Serum creatinine and methotrexate levels should be determined at 24-hour intervals. If the 24-hour serum creatinine has increased 50% over baseline or if the 24-hour methotrexate level is greater than 5×10^{-6} M or the 48-hour level is greater than 9×10^{-7} M, the dose of leucovorin should be increased to 100 mg/m^2 intravenously every 3 hours until the methotrexate level is less than 10^{-8} M.

Hydration (3 L/day) and urinary alkalization with NaHCO_3 should be employed concomitantly. The bicarbonate dose should be adjusted to maintain the urine pH at 7.0 or greater.

In most cases, leucovorin should not be administered simultaneously with systemic methotrexate, since the therapeutic effect of the antimetabolite may be nullified. However, when methotrexate is administered by intra-arterial (regional perfusion) or intrathecal injection, leucovorin may be given (intramuscularly, intravenously or orally) concomitantly, to offset systemic methotrexate toxicity without abolishing the local activity of the cytotoxic drug.

Treatment of Megaloblastic Anemia:

For treatment of megaloblastic anemia due to folate deficiency, the dose should not exceed 1 mg daily. The duration of therapy depends on the hematologic response to the drug. Oral leucovorin is preferred to parenteral therapy, except where cases of severe vomiting impair drug absorption when administered orally.

Treatment of Advanced Colorectal Cancer:

Leucovorin is administered at 200 mg/m^2 by slow intravenous injection prior to dosing with 370 mg/m^2 fluorouracil by slow intravenous injection, for 5 consecutive days.

This 5-day treatment course may be repeated at 4-week (28 days) intervals, provided that the patient has completely recovered from the toxic effects of the prior treatment course.

In subsequent treatment courses, the dosage of fluorouracil should be adjusted based on patient tolerance of the prior treatment course. The daily dosage of fluorouracil should be reduced by 20% for patients who experienced moderate hematologic or gastrointestinal toxicity in the prior treatment course, and by 30% for patients who experienced severe toxicity. For patients who did not experience toxicity in the prior treatment course, fluorouracil dosage may be increased by 10%. Leucovorin dosages are not adjusted for toxicity.

4.3 Reconstitution

Parenteral Products:

Dilution for Intravenous Infusion: When required for intravenous infusion, Leucovorin Calcium Injection USP may be diluted with 5% Dextrose Injection, 0.9% Sodium Chloride Injection, Lactated Ringer's Injection or Ringer's Injection to give a final concentration of 0.05 mg/mL leucovorin. These dilutions may be stored for 24 hours at room temperature. Due to the possibility of antimicrobial contamination, unused solution should be discarded after that time.

Dilutions with the hypertonic infusion solutions, 10% Dextrose Injection and 5% Dextrose in 0.9% Sodium Chloride Injection, may also be prepared to a final concentration of 0.05 mg/mL of leucovorin. However, these dilutions should be stored for no longer than 8 hours at room temperatures. Unused solution should be discarded after that time.

4.4 Administration

Leucovorin Calcium Injection USP may be administered as received by intramuscular injection or intravenous injection, or it may be diluted for intravenous infusion (see 4.3 Reconstitution, Dilution for Intravenous Infusion).

Dispensing of Pharmacy Bulk Vials: Leucovorin Calcium Injection USP, 10 mg/mL, is supplied as 500 mg in 50 mL of sterile, unpreserved, isotonic solution (see 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING).

The use of Pharmacy Bulk Vials is restricted to hospitals with a recognized intravenous admixture program. The Pharmacy Bulk vial is intended for single puncture, multiple dispensing, and for intravenous use only.

Contents of the Pharmacy Bulk vials should be dispensed within eight hours of the initial entry because of the potential for microbial contamination. Discard any unused portion. The diluted solutions prepared from the Pharmacy Bulk Vial should be used within 24 hours if kept at room temperature, from the time of the initial entry into the Pharmacy Bulk vial. However, the product diluted with hypertonic infusion solutions is restricted to 8 hours if kept at room temperature, from the time of the initial entry into the Pharmacy Bulk Vial.

Pharmacy Bulk Vials contain no preservatives. Care must be taken to minimize the potential for inadvertent introduction of microorganisms during manipulation in the hospital environment.

Warning: As with all parenteral drug products, intravenous admixtures should be inspected visually for clarity, particulate matter, precipitate, discoloration and leakage prior to administration, whenever solution and container permit. Solutions showing haziness, particulate matter, precipitate, discoloration or leakage should not be used.

5 OVERDOSAGE

Leucovorin has very low acute and subchronic toxicity in animals. There has been no experience with overdosage of parenteral leucovorin in humans.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Intramuscular injection Intravenous injection Intravenous infusion	Leucovorin Calcium Injection USP is a sterile solution of leucovorin (as the calcium salt), supplied as 10 mg/mL, in water for injection with sodium chloride 8.5 mg/mL, added for isotonicity.	Contains no preservatives. Sodium hydroxide or hydrochloric acid may be used for pH adjustment.

Leucovorin Calcium Injection USP is available as a 10 mg/mL sterile, unpreserved, isotonic solution of leucovorin (as the calcium salt), in 5 mL (50 mg/5 mL) single-use vials.

Leucovorin Calcium Injection USP is available as a 10 mg/mL sterile, unpreserved, isotonic solution of leucovorin (as the calcium salt), in a 50 mL (500 mg/50 mL) Pharmacy Bulk Vial for intravenous administration only. The Pharmacy Bulk Vial is supplied to hospitals with a recognized intravenous admixture program only.

7 WARNINGS AND PRECAUTIONS

Please see 3 SERIOUS WARNINGS AND PRECAUTIONS BOX.

General

Since leucovorin may enhance the toxicity of fluorouracil, combination therapy consisting of leucovorin and fluorouracil for advanced colorectal cancer should be administered under the supervision of a physician experienced in the use of antimetabolite cancer chemotherapy. Particular care should be taken in the treatment of elderly or debilitated colorectal cancer patients, as these patients may be at increased risk of severe toxicity. Death from severe enterocolitis, diarrhea and dehydration has been reported in elderly patients receiving leucovorin and fluorouracil. Concomitant granulocytopenia and fever were present in some but not all of the patients.

In cases of overdosage of folic acid antagonists, prompt administration of leucovorin calcium is essential; if a period of more than four hours intervenes, the treatment may be ineffective due to the time delay. Monitoring of the serum methotrexate concentration is essential in determining the optimal dose of leucovorin to give and duration of therapy. Delayed methotrexate excretion may be an indication of a

third space fluid accumulation (i.e. ascites, pleural effusion), renal insufficiency, low pH of urine or inadequate hydration. Higher doses of leucovorin or prolonged administration may be indicated in such cases. Leucovorin has no apparent effect on pre-existing methotrexate nephrotoxicity.

Excessive leucovorin doses must be avoided since this might impair the antitumour activity of methotrexate, especially in CNS tumours where leucovorin accumulates after repeated courses.

Resistance to methotrexate as a result of decreased membrane transport also implies resistance to folinic acid rescue as both medicinal products share the same transport system.

In case of co-administration of leucovorin and fluorouracil, diarrhea and/or stomatitis/mucositis are the first indications that severe and potentially life-threatening toxicity could develop. Patients who experience these symptoms while receiving any combination therapy regimen involving leucovorin and fluorouracil should be carefully monitored. Generally these symptoms are controllable by reducing the dose of fluorouracil. Treatment-related deaths have been sporadically reported in patients receiving leucovorin/fluorouracil combination therapy.

Leucovorin enhances the toxicity of fluorouracil. When these drugs are administered concurrently in the palliative therapy of advanced colorectal cancer, the dosage of fluorouracil must be reduced. Although the toxicities observed in patients treated with the combination of leucovorin and fluorouracil are qualitatively similar to those observed in patients treated with fluorouracil alone, gastrointestinal toxicities (particularly stomatitis and diarrhea) are observed more commonly and may be more severe in patients receiving the combination therapy.

Leucovorin should be used with caution after methotrexate chemotherapy, when the following medical problems exist:

- Aciduria (urine pH less than 7)
- Ascites
- Dehydration (**Note:** Inadequate hydration including that secondary to vomiting may also result in increased methotrexate toxicity.)
- Gastrointestinal obstruction
- Pleural or peritoneal effusions
- Renal function impairment (**Note:** Risk of methotrexate toxicity is increased because elimination of methotrexate may be impaired and accumulation may occur; even small doses of methotrexate may lead to severe myelosuppression and mucositis; larger doses and/or increased duration of leucovorin treatment may be necessary.)

Patient monitoring is recommended when leucovorin is administered as part of methotrexate chemotherapy programs. Monitoring may include creatinine clearance determinations prior to therapy; plasma or serum methotrexate determinations to detect developing renal function impairment (an increase of greater than 50% within 24 hours is associated with severe renal toxicity); urine pH determination (recommended every 6 hours to ensure that the pH remains greater than 7.0 to minimize the risk of methotrexate nephropathy). Leucovorin has no apparent effect on pre-existing methotrexate nephrotoxicity.

Gastrointestinal

Therapy with leucovorin/fluorouracil must not be initiated or continued in patients who have symptoms of gastrointestinal toxicity of any severity, until those symptoms have resolved. Patients with diarrhea must be closely monitored until the diarrhea has resolved, as rapid clinical deterioration leading to death can occur. Elderly or debilitated patients are at greater risk for severe toxicity when receiving this therapy. In elderly patients, it is recommended to begin with a reduced dosage of fluorouracil.

Monitoring and Laboratory Tests

Calcium levels should be monitored in patients receiving combined leucovorin /5-fluorouracil treatment and calcium supplementation should be provided if calcium levels are low. Complete blood count (CBC) with differential and platelets: prior to each treatment; weekly during the first two courses; at time of anticipated white blood cell (WBC) nadir in all courses thereafter. Electrolytes and liver function tests: prior to each treatment for the first three courses and prior to every other course thereafter.

Reproductive Health: Female and Male Potential

- **Fertility:**

Calcium folinate is an intermediate product in the metabolism of folic acid and occurs naturally in the body. No fertility studies have been conducted with calcium folinate in animals.

7.1 Special Populations

7.1.1 Pregnant Women

Leucovorin is an intermediate product in the metabolism of folic acid and occurs naturally in the body.

There are no adequate and well-controlled clinical studies conducted in pregnant or breast-feeding women.

7.1.2 Breast-feeding

Problems have not been documented. It is unknown if Leucovorin Calcium Injection USP is excreted in human milk. Precaution should be exercised because many drugs can be excreted in human milk.

Calcium folinate in combination with 5-fluorouracil is not recommended for use in woman who are breast-feeding.

7.1.3 Pediatrics

No data are available to Health Canada; therefore, Health Canada has not authorized an indication for pediatric use. Leucovorin may increase the frequency of seizures in susceptible children.

7.1.4 Geriatrics

No information is available regarding the use of leucovorin in geriatrics. Elderly patients are at greater risk of developing severe toxicity when treated with the combination of leucovorin plus fluorouracil for the palliative treatment of colorectal cancer.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

Table 2: Adverse Drug Reactions during Leucovorin Calcium Monotherapy

System Organ Class	ADR Term
General disorders and administrations site conditions	Pyrexia
Immune system disorders	Hypersensitivity
	Anaphylactic reaction
	Anaphylactic shock
Nervous system disorders	Seizure
	Syncope
Skin and subcutaneous tissue disorders	Urticaria
	Stevens-Johnson Syndrome
	Toxic Epidermal Necrolysis

Generally, the safety profile depends on the applied regimen of 5-fluorouracil due to enhancement of the 5-fluorouracil induced toxicities. Additional undesirable effects, when used in combination with 5-fluorouracil, are presented in **Table 3**.

Table 3: Adverse Drug Reactions during Leucovorin Calcium Combination Therapy with 5-fluorouracil

System Organ Class	Adverse Drug Reaction
Blood and lymphatic system disorders	Anemia
	Neutropenia
	Leucopenia
	Thrombocytopenia
Gastrointestinal disorders	Diarrhea
	Nausea
	Vomiting
	Stomatitis
General disorders and administration site conditions	Mucosal inflammation
Metabolism and nutrition disorders	Hyperammonemia
Skin and subcutaneous tissue disorders	Palmar-plantar erythrodysesthesia syndrome (hand-foot syndrome)

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to

the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

Allergic reactions, wheezing, skin rash, hives or itching occur rarely. In combination regimens, the toxicity of fluorouracil is enhanced by leucovorin. The most common manifestations are mucositis, stomatitis, leukopenia and/or diarrhea which may be dose-limiting. In clinical trials with this drug combination, these toxicities were found to be controllable by appropriately reducing the dose of fluorouracil.

8.5 Post-Market Adverse Reactions

Cases of Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN), some fatal, have been reported in patients receiving leucovorin in combination with other agents known to be associated with these disorders. A contributory role of leucovorin in these occurrences of SJS/TEN cannot be excluded.

Fatalities have occurred as a result of gastrointestinal toxicity (predominantly mucositis and diarrhea) and myelosuppression. In patients with diarrhea, rapid clinical deterioration leading to death can occur.

9 DRUG INTERACTIONS

9.1 Serious Drug Interactions

Serious Drug Interactions

- **Patients receiving any combination therapy regimen involving leucovorin and fluorouracil should be carefully monitored for diarrhea and/or stomatitis/mucositis as these are the first indications that severe and potentially life-threatening toxicity could develop (see 7 WARNINGS AND PRECAUTIONS and 9 DRUG INTERACTIONS).**
- **Leucovorin may diminish the effect of anti-epileptic substances such as phenobarbital, primidone and phenytoin. During leucovorin administration in epileptic patients treated with these substances, there is a risk to increase the frequency of seizures due to a decrease of plasma concentrations of anti-epileptic drugs (see 9 DRUG INTERACTIONS).**
- **Leucovorin enhances the cytotoxicity and toxicity of fluorouracil. Leucovorin must not be mixed with fluorouracil in the same intravenous injection or infusion (see 9 DRUG INTERACTIONS).**

9.4 Drug-Drug Interactions

The following drugs or combinations containing these drugs may interact with leucovorin with clinical significance:

- Anticonvulsants, barbiturate
- Anticonvulsant, hydantoin
- Primidone

Large doses of leucovorin may counteract the anticonvulsant effects of these medications.

Leucovorin has been administered simultaneously with pyrimethamine without interfering with its anti-malarial therapy.

Table 4 - Established or Potential Drug-Drug Interactions

Leucovorin Calcium	Source of Evidence	Effect	Clinical comment
anti-epileptic substances such as phenobarbital, primidone and phenytoin	Unknown	diminished	Leucovorin may diminish the effect of anti-epileptic substances such as phenobarbital, primidone and phenytoin. During leucovorin administration in epileptic patients treated with these substances, there is a risk to increase the frequency of seizures (a diminution of plasma levels of enzymatic inductor anticonvulsant drugs may be observed because the hepatic metabolism is increased, as folates are one of the cofactors). Clinical monitoring, possibly monitoring of the plasma concentrations and, if necessary, dose adaptation of the anti-epileptic drug during leucovorin administration and after discontinuation is recommended.
folic acid antagonist (e.g., cotrimoxazole, pyrimethamine, methotrexate, antibiotic with antifolic effect)	Unknown	Reduced or neutralized	When leucovorin is given in conjunction with a folic acid antagonist (e.g., cotrimoxazole, pyrimethamine, methotrexate, antibiotic with antifolic effect), the efficacy of the folic acid antagonist may either be reduced or completely neutralized.
Di-aminopyrimidines (e.g., trimethoprim or co-trimoxazole)	Unknown	Inhibit antibiotic effect	Di-aminopyrimidines (there is some evidence that concomitant administration of leucovorin and trimethoprim (or co-trimoxazole) may inhibit the antibiotic effect of trimethoprim)
Methotrexate	unknown	Nullify antitumour effect	Leucovorin administered concomitantly with methotrexate may nullify the antitumour chemotherapeutic effect of the latter drug (see 4 DOSAGE AND ADMINISTRATION).

Leucovorin Calcium	Source of Evidence	Effect	Clinical comment
Fluorouracil	Unknown	Enhance cytotoxicity	Leucovorin enhances the cytotoxicity and toxicity of fluorouracil. Leucovorin must not be mixed with fluorouracil in the same intravenous injection or infusion.

9.5 Drug-Food Interactions

Interactions with food have not been established.

9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.7 Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been established.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Leucovorin is a reduced form of folic acid, which is readily converted to other reduced folic acid derivatives (e.g., tetrahydrofolate). Because it does not require reduction by dihydrofolate reductase as does folic acid, leucovorin is not affected by blockages of this enzyme by folic acid antagonists (dihydrofolate reductase inhibitors). This allows purine and thymidine synthesis, and thus DNA, RNA and protein synthesis to occur. Leucovorin may limit methotrexate action on normal cells by competing with methotrexate for the same transport processes into the cell.

Leucovorin enhances the cytotoxicity of fluoropyrimidines such as fluorouracil by their metabolites, methylene tetrahydrofolate and fluorodeoxyuridine monophosphate, forming a stable ternary complex with thymidylate synthase and thereby decreasing intracellular levels of that enzyme and the product thymidylate. The cell then dies as a result of thymine starvation.

10.2 Pharmacodynamics

A folic acid deficiency is produced during therapy with the folic acid antagonists aminopterin and amethopterin (methotrexate) used as antineoplastic agents and with the chemotherapeutic agent pyrimethamine. These agents competitively inhibit the conversion of folic acid to folinic acid. Their affinity for folate reductase is so much greater than that of folic acid that not even large doses of folic acid will correct the drug-induced deficiency. In the event of a severe toxic reaction, the already reduced form, folinic acid, can be given, since it can be used directly to form new coenzyme.

10.3 Pharmacokinetics

Thirty minutes following oral administration of leucovorin calcium, 92 to 93% of total reduced folates in serum were assayable as 5-methyltetrahydrofolate. The determination of 5-methyltetrahydrofolate was carried out by the use of a differential microbiological disc assay procedure utilizing methotrexate

resistant strains of *Lactobacillus casei* and *Streptococcus faecium var. durans*. Peak serum levels of 5-methyltetrahydrofolate were reached earlier following intramuscular administration (approximately 45 minutes) than after oral administration (approximately 2 hours).

Similar results were obtained after oral administration of radiolabelled leucovorin calcium. These studies also indicated substantial metabolism of leucovorin calcium during transfer from gastrointestinal tract to the systemic circulation, since 90% of the serum folate was identified as 5-methyltetrahydrofolate by chromatographic techniques.

The serum half-life of leucovorin (or 5-formyltetrahydrofolate) was 35 to 45 minutes following both oral and intramuscular administration. The serum half-life of 5-methyltetrahydrofolate was about 2 1/4 hours. 5-methyltetrahydrofolate was excreted via the kidneys in a manner proportional to its serum concentration.

Methotrexate did not seem to affect the absorption of folate.

11 STORAGE, STABILITY AND DISPOSAL

Leucovorin Calcium Injection USP should be stored refrigerated (2°C to 8°C) and protected from light and freezing.

12 SPECIAL HANDLING INSTRUCTIONS

There are no special handling instructions for this drug product.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: Leucovorin calcium. Also known as calcium folinate, citrovorum factor and folinic acid.

Chemical name: L-glutamic acid, N (4 ((2 amino 5 formyl 1, 4, 5, 6, 7, 8 hexahydro 4 oxo 6 pteridiny)methyl)amino)benzoyl) calcium salt (1:1).

Molecular formula and molecular mass: $C_{20}H_{21}CaN_7O_7 \cdot 5H_2O$, 601.62

Structural formula:



Physicochemical properties: Leucovorin is a mixture of the diastereoisomers of 5-formyltetrahydrofolic acid, the biologically active component of the mixture being the (-)-L-isomer.

Leucovorin calcium occurs as a yellowish to beige, odourless powder, which is very soluble in water and practically insoluble in alcohol. The hydrated salt contains 8 to 15% water. The optical rotation (concentration = 2.5% in water) is $+14.5^\circ$ to $+16.5^\circ$ for the anhydrous substance. The pH is 7.5 to 8.5 for this 2.5% aqueous solution.

14 CLINICAL TRIALS

The clinical trial data on which the original indication was authorized are not available.

15 MICROBIOLOGY

No microbiological information is required for this drug product.

16 NON-CLINICAL TOXICOLOGY

In mice, the LD_{50} was 991 mg/kg intravenously. Toxic symptoms included body tremors, marked ataxia, clonic convulsions and deaths within 10 minutes in CD-1 male mice. The single-dose oral LD_{50} could not be determined because, at doses as high as 20,000 mg/kg, no toxicity was observed in CD-1 and Long-Evans male rats. Doses higher than this could not be given because of the limitations of dose volume and viscosity.

In subchronic studies, oral doses of leucovorin at 0, 75, 225 or 675 mg/kg daily for over 30 days to rats and beagle dogs produced no drug-related toxic effects on body weight, food consumption, hematology, blood chemistry, urinalysis or pathology. No alteration in ECG in dogs occurred. Ophthalmoscopic examinations of rats and dogs revealed no drug-induced toxic effects.

Genotoxicity, carcinogenicity, and fertility studies have not been conducted with calcium folinate.

Embryo-fetal reproduction toxicity studies have been performed in rats and rabbits. Rats were dosed up to 1800 mg/m² which is 9 times the maximum recommended human dose, and rabbits were dosed up to 3300 mg/m² which is 16 times the maximum recommended human dose. There was no embryo-fetal toxicity noted in rabbits. At the maximum dose in rats, there was a slight increase in early embryonic resorptions and no other adverse effects on embryo-fetal development. No resorptions were noted in dose groups at 5 times the maximum recommended human dose.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

^{Pf}LEUCOVORIN CALCIUM INJECTION USP

Leucovorin calcium

Read this carefully before you receive **Leucovorin Calcium Injection USP**. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Leucovorin Calcium Injection USP**.

Serious Warnings and Precautions

- You will receive Leucovorin Calcium Injection USP:
 - as a shot into your muscle (intramuscularly) or as a shot or infusion (drip) into your vein (intravenously). It must not be given into the fluid-filled space between the thin layers of tissue that cover the brain and spinal cord (intrathecally). Deaths have happened in people getting this medicine into the spine.
 - with 5-fluorouracil or methotrexate only under the care of a healthcare professional experienced in the use of anti-cancer medicines.
- If you are receiving leucovorin with fluorouracil (an anticancer medicine), your healthcare professional will monitor you for side effects like diarrhea or painful sores that may make it hard for you to eat, drink, or swallow (stomatitis/mucositis). These are early signs of severe and potentially life-threatening side effects. Talk to your healthcare professional **right away** if you experience any of these side effects.
- Deaths from the following side effects have happened in people getting leucovorin:
 - **Severe problems with the digestive system** (gastrointestinal toxicity): These include diarrhea and mucositis (ulcers (open sores) and swelling of lining of the bowels and mouth).
 - **Severe drop in blood cell counts** (myelosuppression, a condition in which the bone marrow cannot make enough blood cells).
 - **Severe skin reactions**: These include Stevens-Johnson syndrome (SJS) and Toxic Epidermal Necrolysis (TEN). These are more likely to occur if you are taking other medicines known to cause these skin reactions.For more information on these and other serious side effects, see the **Serious side effects and what to do about them** table, below.
- Leucovorin may lower the effects of anti-epileptic medications such as phenobarbital, primidone and phenytoin. If you are taking anti-epileptic medications, you may be at an increased risk of having seizures more often

What is Leucovorin Calcium Injection USP used for?

Leucovorin Calcium Injection USP is used:

- to lower the toxic effects of:
 - a group of medicines called folic acid antagonists. It is also used to treat an overdose of these medicines
 - methotrexate (a medicine often used to treat cancer)
- to treat some kinds of anemia (low level of red blood cells in the body)
 - due to low level of folate (an essential B vitamin) in the body such as in poor food intake or absorption (sprue, nutritional deficiency); or
 - occurring during pregnancy and infancy.
- before treatment with fluorouracil to improve survival time in the palliative treatment of patients with advanced colorectal cancer.

How does Leucovorin Calcium Injection USP work?

Leucovorin Calcium Injection USP belongs to a class of medicines called folic acid analogs. It:

- lowers the harmful effects of methotrexate by competing with methotrexate and limiting how much methotrexate enters your cells
- treats anemia by supplying folic acid that is needed to produce red blood cell
- increases the effects of 5-fluorouracil (a medicine used to treat cancer)

What are the ingredients in Leucovorin Calcium Injection USP?

Medicinal ingredients: Leucovorin Calcium (also known as calcium folinate).

Non-medicinal ingredients: Hydrochloric acid or sodium hydroxide (to adjust pH).

Leucovorin Calcium Injection USP comes in the following dosage forms:

Solution: 10 mg / mL

Do not receive Leucovorin Calcium Injection USP if:

- you have a type of anemia caused by too little Vitamin B₁₂
- you are allergic (hypersensitive) to calcium folinate or any of the other ingredients of Leucovorin Calcium Injection USP (See What are the ingredients in Leucovorin Calcium Injection USP)?

Leucovorin Calcium Injection USP must NOT be injected into the fluid-filled space between the thin layers of tissue that cover the brain and spinal cord (intrathecally).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you receive Leucovorin Calcium Injection USP. Talk about any health conditions or problems you may have, including if you:

- are currently taking methotrexate and:
 - have aciduria (urine pH less than 7)
 - have or have ever had a build-up of fluid in the chest cavity or the belly area
 - are dehydrated (feeling thirsty, dark yellow urine, dizzy, lightheaded, tired, dry mouth and peeing less than 4 times a day)

- have stomach or bowel issues such as a blockage
- you have trouble with your kidneys
- have diarrhoea
- are pregnant, plan to become pregnant, or are breast-feeding.

Other warnings you should know about:

- **Older people with colorectal cancer.** You could have more side effects. Deaths from severe bowel problems, diarrhea, and dehydration have happened in older people getting leucovorin with fluorouracil.
- **Lab and blood tests:** Your healthcare professional will do blood tests before you receive Leucovorin Calcium Injection USP and/or during treatment to monitor your progress or check for side effects. These tests may check:
 - the level of blood cells in your body.
 - that your liver or kidneys are working properly.
 - the level calcium and other essential minerals—like sodium and potassium (electrolyte) in your blood.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

- If you are receiving leucovorin with fluorouracil (an anticancer medicine):
 - your healthcare professional will monitor you for side effects like diarrhea or painful sores that may make it hard for you to eat, drink, or swallow (stomatitis/mucositis). These are early signs of severe and potentially life-threatening side effects. Talk to your healthcare professional **right away** if you experience any of these side effects. See the **Serious side effects and what to do about them** table, below
 - you may experience increased side effects from fluorouracil
- Leucovorin may lower the effects of anti-epileptic medications such as phenobarbital, primidone and phenytoin. If you are taking anti-epileptic medications you may be at an increased risk of having seizures more often
- Leucovorin must not be mixed in the same infusion as 5-fluorouracil because a precipitate may form

The following may interact with Leucovorin Calcium Injection USP:

- medicines used to treat epilepsy - phenobarbital, primidone, and phenytoin. Your healthcare professional may check blood levels of these medicines and change your dose to prevent increased convulsions (fits).
- a group of medicines called folic acid antagonists). These include:
 - methotrexate (a medicine often used to treat cancer). High doses of leucovorin may lower how well methotrexate works
 - trimethoprim or co-trimoxazole (an antibiotic)
 - pyrimethamine (a medicine used to treat malaria)

- anti-cancer medicines, such as 5-fluorouracil (5FU). Leucovorin may increase the harmful effect of fluorouracil

How you will receive Leucovorin Calcium Injection USP:

- in a healthcare setting under the care of your healthcare professional
- as an injection into the muscle or as an injection or an infusion (drip) into your vein

Usual dose:

Your healthcare professional will work out the right dose of Leucovorin Calcium Injection USP for you.

Overdose:

If you think you, or a person you are caring for, have received too much Leucovorin Calcium Injection USP, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

What are possible side effects from using Leucovorin Calcium Injection USP?

These are not all the possible side effects you may have when taking Leucovorin Calcium Injection USP. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- nausea, vomiting
- red, swollen lips
- dizziness
- fever

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
VERY COMMON			
Gastrointestinal toxicity (Severe problems with the digestive system): <ul style="list-style-type: none"> • Diarrhea: persistent or severe • Stomatitis (swelling and redness of the lining of your mouth): painful sores that may make it hard for you to eat, drink, or swallow • Mucositis (ulcers (open sores) and swelling of lining of the bowels and mouth): red, shiny, 			

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
or swollen mouth and gums, sores in the mouth or on the gums or tongue, blood in the mouth, pain in the mouth or throat, difficulty swallowing or talking, mild burning, or pain when eating			
COMMON			
Palmar-plantar erythrodysesthesia (Hand and Foot syndrome): red or swollen palms, thick calluses and blisters of the hands and soles of the feet, tingling or burning, tightness of the skin		✓	
UNKNOWN			
Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) (severe skin reactions): redness, blistering and/or peeling of the skin and/or inside the lips, mouth, eyes, nasal passages or genitals, accompanied by fever, chills, tiredness, headache and cough, body aches or swollen glands, raised red or purple skin patches, possibly with blister or crust in the center, swollen lips, mild itching or burning			✓
Allergic reactions: difficulty swallowing or breathing, wheezing, drop in blood pressure, feeling sick to your stomach and throwing up, hives or rash, swelling of the face, lips, tongue or throat, low blood pressure, confusion, reduced alertness, cold, moist skin, fast breathing, fast heartbeat			✓
Seizures (fit): uncontrollable shaking with or without loss of consciousness			✓
Bone marrow suppression (severe drop in blood cells): bleeding,			✓

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
bruising, chills, fatigue, fever, infections, weakness, shortness of breath or other signs of infection			
Syncope (fainting): a temporary loss of consciousness due to a sudden drop in blood pressure		✓	
Hyperammonemia (high ammonia levels in the blood): confusion, irritability, refusal to eat meat or high protein products		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Leucovorin Calcium Injection USP will be stored by your healthcare professional. It should be kept:

- refrigerated (2°C to 8°C)
- protected from light and freezing
- out of reach and sight of children.

If you want more information about Leucovorin Calcium Injection USP:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: <https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug->

[product-database.html](#); the manufacturer's website www.Pfizer.ca, or by calling 1-800-463-6001.

This leaflet was prepared by Pfizer Canada ULC.

Last Revised DEC 13, 2022