

PART III: CONSUMER INFORMATION

VIAGRA[®]
(sildenafil citrate tablets)

This leaflet is part III of a three-part "Product Monograph" published when VIAGRA was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about VIAGRA. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

VIAGRA[®] (sildenafil citrate) is a treatment for erectile dysfunction in male adults. This is when a man cannot achieve or maintain a hard, erected penis for satisfactory sexual activity.

It is important to note that VIAGRA works only with sexual stimulation. VIAGRA alone does not increase your sex drive.

What it does:

VIAGRA belongs to a class of medicine called phosphodiesterase type 5 (PDE5) inhibitors. Following sexual stimulation, VIAGRA works by helping to relax the blood vessels in your penis by allowing blood to flow into your penis. This results in improved erectile function.

When it should not be used:

- If you are taking any medicines containing nitrates in any form (oral, sublingual [under the tongue], skin patch, or by inhalation [spray]). Although VIAGRA is used occasionally, nitrates must never be used. Nitrates are found in many prescription medicines that are used in the treatment of angina pectoris (chest pain due to heart disease), such as nitroglycerin, isosorbide mononitrate, or isosorbide dinitrate. If you do not understand what nitrates are, or are unsure about whether a medication you are taking is a "nitrate", ask your doctor or pharmacist.

If you take VIAGRA with nitrate-containing medicines or any other nitrates (e.g., amyl nitrite "poppers"), your blood pressure could suddenly drop to a life-threatening level. You could get dizzy, faint, or even have a heart attack or stroke.

- If you have loss of vision in one or both eyes from an eye disease called non-arteritic anterior ischaemic optic neuropathy (NAION)
- If you have ever had an allergic reaction to sildenafil citrate or the nonmedicinal ingredients in VIAGRA.
- VIAGRA is not to be used in men for whom sexual activity is inadvisable.
- Do not take VIAGRA with guanylate cyclase stimulators, such as riociguat.

What the medicinal ingredient is:

sildenafil citrate.

What the nonmedicinal ingredients are:

microcrystalline cellulose, anhydrous dibasic calcium phosphate, croscarmellose sodium, magnesium stearate, hypromellose, titanium dioxide, lactose, triacetin, and FD & C Blue #2 aluminum lake.

What dosage forms it comes in:

VIAGRA is available as blue, rounded, diamond-shaped tablets marked 'PFIZER' on one side, and 'VGR 25, 50, or 100' on the other. Each tablet contains either 25 mg, 50 mg or 100 mg sildenafil citrate.

WARNINGS AND PRECAUTIONS

This medicine has been prescribed for you personally and you should not pass it on to others. It may harm them, even if their symptoms are the same as yours.

BEFORE you use VIAGRA talk to your doctor or pharmacist if you:

- have heart problems (irregular heart beats, angina, chest pain, or had a heart attack). If you have heart problems, ask your doctor if your heart is healthy enough to handle the extra strain of having sex. If you have chest pain, dizziness or nausea during sex, stop exerting yourself. You should **not** use nitrates but you should seek immediate medical assistance.
- are 65 years of age or over
- have had a stroke
- have low blood pressure or uncontrolled high blood pressure
- have liver or kidney problems
- have sickle cell anemia (abnormality of the red blood cells), multiple myeloma (cancer of the bone marrow) or leukaemia (cancer of the white blood cells)
- are allergic to sildenafil or any of the other ingredients of VIAGRA tablets
- have a deformed penis or Peyronie's disease
- have ever had an erection that lasted more than 4 hours
- have stomach ulcers or other bleeding problems.
- have a rare inherited eye disease called retinitis pigmentosa
- have had temporary, decrease, or permanent loss of vision in one or both eyes, including a condition called Non-Arteritic Anterior Ischemic Optic Neuropathy (NAION). The specific type of vision decrease or loss, called non-arteritic anterior ischemic optic neuropathy (NAION), seems to occur rarely when blood flow to the optic nerve is reduced or blocked. Vision decrease or loss may be partial or complete, in one or very occasionally both eyes. While in some cases the condition may improve over time, it can also be irreversible.
- Rare but serious skin reactions of Stevens-Johnson's Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN) have been reported during the post-marketing period.

VIAGRA is not recommended for use in children under 18 years of age and in women.

VIAGRA does not protect against sexually transmitted diseases (STD), including Human Immunodeficiency Virus (HIV).

Alcohol consumption may decrease the ability to get an erection.

If you are taking **VIAGRA** and experience temporary, decrease, or permanent loss of vision, stop taking **VIAGRA** and call your doctor.

In case of chest pain occurring during or after sexual activity you should **not** use nitrates but you should seek immediate medical assistance.

Sudden decrease or loss of hearing has been reported in a few number of postmarketing and clinical trial cases with the use of PDE5 inhibitors, including **VIAGRA**. It has not been established whether these are related directly to the use of these medications or to other factors. If you experience these symptoms, stop taking **VIAGRA** and call your doctor.

Driving and using machines: Before you perform tasks which may require special attention, wait until you know how you respond to **VIAGRA**. Dizziness or altered vision can occur while using **VIAGRA**.

INTERACTIONS WITH THIS MEDICATION

You should tell your doctor or pharmacist about any other medications that you are taking. **VIAGRA** may interfere with some drugs. Only take **VIAGRA** with other drugs if recommended by your doctor.

Drugs that may interact with **VIAGRA** include:

- any drugs that contain nitrates in any form (oral, sublingual [under the tongue], skin patch or by inhalation [spray]). Nitrates are found in many prescriptions that are used to treat angina pectoris (chest pain due to heart disease). You should not take **VIAGRA** if you are taking these drugs.
- alpha-blockers (drugs used to treat prostate problems or high blood pressure)
- ketoconazole or itraconazole (drugs used to treat fungal infections)
- erythromycin (a drug used to treat bacterial infections)
- ritonavir, saquinavir or other drugs for the treatment of HIV
- cimetidine (a drug generally used to treat duodenal or gastric problems)
- bosentan (a drug used in the treatment of high blood pressure in the blood vessels between the heart and the lungs)
- drugs to treat erectile dysfunction. These drugs are Cialis, Levitra, Adcirca and Staxyn.

- other drugs that contain sildenafil. They can be called Revatio. They are used in the treatment of high blood pressure in the blood vessels between the heart and the lungs.

Grapefruit juice may increase the levels of **VIAGRA** in your blood.

PROPER USE OF THIS MEDICATION

Usual dose:

Your doctor can determine the dose that works best for you. Always take **VIAGRA** as prescribed by your doctor. You should speak with your doctor or pharmacist if you are unsure. A dose above 100 mg per day is not recommended.

Swallow the tablet whole, with some water.

You should **not** take more than one dose of **VIAGRA** per day.

If you have serious liver or kidney problems or you are 65 years of age or over, your doctor may start you at the lowest dose (25 mg) of **VIAGRA**.

How to optimize your use of **VIAGRA**:

You should take **VIAGRA** approximately 30 to 60 minutes before sexual activity. You can engage in sexual activity within 30 minutes of taking the tablet and for up to 4 hours. The amount of time it takes to have an effect varies slightly from person to person. Remember, sexual stimulation is required for **VIAGRA** to work.

You should avoid excessive drinking of alcohol, since alcohol can temporarily impair the ability to get an erection.

VIAGRA can be taken with or without food. However, you may find that it takes longer for **VIAGRA** to work if you take it with a high-fat meal.

VIAGRA may not work the first time or every time. If **VIAGRA** did not work for you on one occasion, try again on another day. You will learn how well **VIAGRA** works for you through your personal experience. The first few times may be charged with emotion or anxiety. If after a few separate attempts, you do not get the results you expect, talk to your doctor.

Overdose:

You should **not** take more than one dose of **VIAGRA** per day. If you have taken more Viagra than you should, contact your doctor or a poison control centre immediately.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

As with most drugs, **VIAGRA** can cause some side effects. These effects are usually mild to moderate in nature and do not last for a long time.

Side effects may include:

- headache, facial flushing

- nausea, vomiting, indigestion, abdominal pain,
- dizziness
- dry, stuffy, or swollen nose
- throat tightness, dry mouth, decreased sensitivity of the mouth
- pain in arms and legs, myalgia (muscle pain)
- somnolence
- erection increased

If you notice any side effects not mentioned above, or any of the above-mentioned side effects persist or become bothersome, please contact your doctor or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

	Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Less common	Effect on hearing: sudden decrease or loss of hearing		✓	
	Effects on vision: colour tinge, increased brightness of light, blurred vision			✓
	Impaired or sudden loss of vision: decreased eyesight or unable to see with one or both eyes			✓
	Detached retina: a decrease in, or sudden loss of vision in one or both eyes			✓
Rare	Serious skin reactions: rash, blisters, peeling skin and pains.			✓
Very rare	Priapism: erection lasting more than 4 hours			✓
Unknown	Cough		✓	
	Allergic reactions: rash, hives, itch, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			✓
	Nosebleed		✓	

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
<i>Noted in patients taking sildenafil citrate for pulmonary hypertension:</i> Fever		✓	
Shortness of breath		✓	
Seizure, seizure recurrence			✓
Transient global amnesia: temporary memory loss		✓	
Heart attack (myocardial infarction): chest pain, shortness of breath			✓
Chest pain (unstable angina)			✓
Arrhythmia/tachycardia, palpitations: fast or irregular heart beat, heart rate increased			✓
Hypotension (low blood pressure): dizziness, fainting, lightheadedness	✓		
Stroke (cerebrovascular hemorrhage): bleeding in the brain, vision changes, difficulty speaking, weakness on one side of the body, dizziness, lack of coordination or poor balance			✓
Transient ischaemic attack: temporary loss of vision, difficulty speaking, weakness on one side of the body, numbness or tingling usually on one side of the body, dizziness, lack of coordination or poor balance.			✓

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Pulmonary Hemorrhage (acute bleeding from the lung): oozing of bloody fluid from the nose and respiratory tract, accompanied by rapid worsening of patient respiration, turning blue and in severe cases, shock)			✓

This is not a complete list of side effects. For any unexpected effects while taking VIAGRA, contact your doctor or pharmacist.

HOW TO STORE IT

- Store between 15°C and 30°C, in the original package. Do not freeze.
- Do not take VIAGRA after the expiry date shown on the package.
- Always keep VIAGRA out of reach and sight of children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: **Canada Vigilance Program
Health Canada
Postal Locator 1908C
Ottawa, Ontario
K1A 0K9**

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at: <http://www.pfizer.ca> or by contacting the sponsor, Pfizer Canada Inc., at: 1-800-463-6001

This leaflet was prepared by Pfizer Canada Inc.

Last revised: 18 September 2018