

PRODUCT MONOGRAPH

PrGD[®]-diclofenac/misoprostol 50

diclofenac sodium and misoprostol enteric-coated tablets

50 mg diclofenac/200 µg misoprostol

PrGD[®]-diclofenac/misoprostol 75

diclofenac sodium and misoprostol enteric-coated tablets

75 mg diclofenac/200 µg misoprostol

NSAID with a
Mucosal Protective Agent

GenMed, a division of Pfizer Canada ULC
17,300 Trans-Canada Highway
Kirkland, Quebec H9J 2M5
www.pfizer.ca

Date of Preparation:
January 21, 2010
Date of Revision:
November 2, 2020

Submission Control No: 244809

[®] GD is a trademark of Pfizer Canada ULC
GenMed, a division of Pfizer Canada ULC, Licensee
© Pfizer Canada ULC 2020

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....3

SUMMARY PRODUCT INFORMATION3

INDICATIONS AND CLINICAL USE.....3

CONTRAINDICATIONS4

WARNINGS AND PRECAUTIONS.....6

ADVERSE REACTIONS.....17

DRUG INTERACTIONS22

DOSAGE AND ADMINISTRATION26

OVERDOSAGE27

ACTION AND CLINICAL PHARMACOLOGY27

STORAGE AND STABILITY30

DOSAGE FORMS, COMPOSITION AND PACKAGING30

PART II: SCIENTIFIC INFORMATION31

PHARMACEUTICAL INFORMATION.....31

CLINICAL TRIALS.....32

DETAILED PHARMACOLOGY33

TOXICOLOGY34

REFERENCES42

PART III: CONSUMER INFORMATION.....45

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

PrGD-diclofenac/misoprostol 50

PrGD-diclofenac/misoprostol 75

diclofenac sodium and misoprostol enteric-coated tablets

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Enteric coated tablets: 50 mg diclofenac/ 200 µg misoprostol 75 mg diclofenac /200 µg misoprostol	Lactose <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol) is indicated for the following:

- Acute and chronic use in the relief of the signs and symptoms of rheumatoid arthritis and osteoarthritis.

Throughout this document, the term NSAIDs refers to both non-selective NSAIDs and selective COX-2 inhibitor NSAIDs, unless otherwise indicated.

Diclofenac, particularly at higher doses, is associated with an increased risk of serious cardiovascular related adverse events that is comparable to COX-2 inhibitors and high dose ibuprofen. For patients with pre-existing risk factors for cardiovascular disease (including ischemic heart disease, cerebrovascular disease and/or congestive heart failure NYHA II-IV) other management strategies that do not include NSAIDs, particularly COX-2 inhibitors, ibuprofen and diclofenac, should be considered first (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

For patients with increased risk of developing GI adverse events other management strategies that do not include NSAIDs should be considered first (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

Use of GD-diclofenac/misoprostol should be limited to the lowest effective dose for the shortest possible duration of treatment in order to minimize the potential risk for cardiovascular or gastrointestinal adverse events (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS).

GD-diclofenac/misoprostol, as a NSAID, does NOT treat clinical disease or prevent its progression.

GD-diclofenac/misoprostol, as a NSAID, only relieves symptoms and decreases inflammation for as long as the patient continues to take it.

Geriatrics (>65 years of age):

Evidence from clinical studies and postmarket experience suggests that use in geriatric patients is associated with differences in safety or effectiveness (see **WARNINGS AND PRECAUTIONS**).

Pediatrics (<18 years of age):

Not recommended for pediatric use (see **CONTRAINDICATIONS**).

CONTRAINDICATIONS

GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol) is contraindicated in:

- The peri-operative setting of coronary artery bypass graft surgery (CABG). Although GD-diclofenac/misoprostol has NOT been studied in this patient population, a selective COX-2 inhibitor NSAID studied in such a setting has led to an increased incidence of cardiovascular/thromboembolic events, deep surgical infections and sternal wound complications.
- The women who are **pregnant**, or in whom **pregnancy has not been excluded**. Women of childbearing potential should be fully counseled about misoprostol’s abortifacient potential and the importance of effective contraception (oral contraceptive or intrauterine device) and prevention of pregnancy while undergoing treatment (see **WARNINGS AND PRECAUTIONS – Special Populations – *Pregnant Women***).
- Women who are breastfeeding, because of the potential for serious adverse reactions in nursing infants.
- Severe uncontrolled heart failure.
- Known or suspected hypersensitivity to diclofenac sodium, misoprostol, or any of the components/excipients.
- History of asthma, urticaria, or allergic-type reactions after taking ASA or other NSAIDs (i.e. complete or partial syndrome of ASA-intolerance - rhinosinusitis, urticaria/angioedema, nasal polyps, asthma). Fatal anaphylactoid reactions have occurred in such individuals. As well, individuals with the above medical problems are at risk of a severe reaction even if they have taken NSAIDs in the past without any adverse reaction. The potential for cross-reactivity between different NSAIDs must be kept in mind (see **WARNINGS AND PRECAUTIONS – Hypersensitivity Reactions – *Anaphylactoid Reactions***).

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

- Patients with active gastric / duodenal / peptic ulcer, active gastrointestinal bleeding, a history of recurrent ulceration or active inflammatory disease of the gastrointestinal system.
- Cerebrovascular bleeding or other bleeding disorders.
- Inflammatory bowel disease.
- Significant hepatic impairment or active liver disease.
- Severely impaired or deteriorating renal function (creatinine clearance <30 mL/min or 0.5 mL/sec). Individuals with lesser degrees of renal impairment are at risk of deterioration of their renal function when prescribed NSAIDs and must be monitored (see **WARNINGS AND PRECAUTIONS – Renal**).
- Known hyperkalemia (see **WARNINGS AND PRECAUTIONS – Renal – Fluid and Electrolyte Balance**).
- Children and adolescents less than (18) years of age.
- Diclofenac is contraindicated for use with other NSAIDs because of the absence of any evidence demonstrating synergistic benefits and the potential for additive side effects (see **DRUG INTERACTIONS**).

WARNINGS AND PRECAUTIONS

Risk of Cardiovascular (CV) Adverse Events: Cardiovascular Disease (including Ischemic Heart Disease, Cerebrovascular Disease, Congestive Heart Failure (NYHA II-IV) (see WARNINGS AND PRECAUTIONS – Cardiovascular)

Diclofenac is associated with an increased risk of serious cardiovascular (CV) thrombotic events (such as myocardial infarction and stroke), which can be fatal. The increased risk is comparable to COX-2 inhibitors and high-dose ibuprofen. An increased risk of CV serious thrombotic events may occur early in the treatment and become higher with the duration of treatment. The risk may increase with the dose. Patients with CV disease or CV risk factors may be at greater risk (See Cardiovascular in WARNINGS AND PRECAUTIONS).

For patients with a high risk of developing an adverse CV event, other management strategies that do NOT include NSAIDs, particularly COX-2 inhibitors, ibuprofen and diclofenac, should be considered first. **To minimize the potential risk for an adverse CV event, the lowest effective dose should be used for the shortest possible duration.**

Treatment with GD-diclofenac/misoprostol is not recommended in patients with pre-existing cardiovascular disease (congestive heart failure NYHA -II-IV, ischemic heart disease, peripheral arterial disease) cerebrovascular disease, uncontrolled hypertension or patients with risk factors for cardiovascular disease (e.g. hypertension, hyperlipidemia, diabetes mellitus and smoking). These patients should be treated with GD-diclofenac/misoprostol only after careful consideration.

Use of NSAIDs, such as GD-diclofenac/misoprostol, can promote sodium retention in a dose-dependent manner, through a renal mechanism, which can result in increased blood pressure and/or exacerbation of congestive heart failure (see also WARNINGS AND PRECAUTIONS – Renal – *Fluid and Electrolyte Balance*).

Risk of Gastrointestinal (GI) Adverse Events (see WARNINGS AND PRECAUTIONS – Gastrointestinal)

Use of NSAIDs, such as GD-diclofenac/misoprostol, is associated with an increased incidence of gastrointestinal adverse events (such as peptic/duodenal ulceration, perforation, obstruction and gastrointestinal bleeding).

General

Frail or debilitated patients may tolerate side effects less well and therefore special care should be taken in treating this population. **To minimize the potential risk for an adverse event, the lowest effective dose should be used for the shortest possible duration.** As with other NSAIDs, caution should be used in the treatment of elderly patients who are more likely to be suffering from impaired renal, hepatic or cardiac function. For high risk patients, alternate therapies that do not involve NSAIDs should be considered.

The use of diclofenac/misoprostol is NOT recommended for use with other NSAIDs, with the

exception of low-dose ASA for cardiovascular prophylaxis, because of the absence of any evidence demonstrating synergistic benefits and the potential for additive adverse reactions including gastrointestinal ulcers and bleeding (see **DRUG INTERACTIONS – Drug-Drug Interactions – Acetylsalicylic acid (ASA) or other NSAIDs**).

In common with other anti-inflammatory drugs, GD-diclofenac/misoprostol may mask the usual signs of infection, such as fever.

Carcinogenesis and Mutagenesis

(See **TOXICOLOGY**)

Cardiovascular

Diclofenac is associated with an increased risk of serious cardiovascular (CV) thrombotic events (such as myocardial infarction and stroke), which can be fatal. The increased risk is comparable to COX-2 inhibitors and high-dose ibuprofen. An increased risk of CV serious thrombotic events may occur early in the treatment and become higher with the duration of treatment. The risk may increase with the dose. Patients with CV disease or CV risk factors may be at greater risk.

Meta-analyses of randomized clinical trials comparing several different NSAIDs suggest that diclofenac, particularly at higher doses, is associated with an increased risk of cardiovascular adverse events that is comparable to COX-2 inhibitors and high dose ibuprofen. Large population-based observational studies conducted in the general population also support these findings. Some observational studies showed that the increased risk of the CV thrombotic events began as early as the first weeks of treatment. Such risk increased with duration of NSAID treatment. The relative increase in risk of serious CV thrombotic events during NSAID treatment appears to be similar in patients with or without CV disease or CV risk factors. However, patients with CV disease or CV risk factors during the treatment had a higher absolute risk of serious CV thrombotic events due to their increased baseline rate.

As the cardiovascular risks of diclofenac may increase with dose and duration of exposure, the lowest effective daily dose should be used for the shortest duration possible. The patient's need for symptomatic relief and response to therapy should be re-evaluated periodically.

Patients should remain alert for the signs and symptoms of serious arteriothrombotic events (e.g. chest pain, shortness of breath, weakness, slurring of speech), which can occur without warnings. Patients should be instructed to see a physician immediately in case of such an event.

Use of NSAIDs, such as GD-diclofenac/misoprostol, can lead to new hypertension or can worsen pre-existing hypertension, either of which may increase the risk of cardiovascular events as described below. Thus blood pressure should be monitored regularly. Consideration should be given to discontinuing GD-diclofenac/misoprostol should hypertension either develop or worsen with its use.

Use of NSAIDs, such as **GD-diclofenac/misoprostol**, can induce fluid retention and edema, and may exacerbate congestive heart failure, through a renally-mediated mechanism (See **WARNINGS AND PRECAUTIONS - Renal - Fluid and Electrolyte Balance**).

Caution should be exercised in prescribing GD-diclofenac/misoprostol to patients with risk factors for cardiovascular disease, cerebrovascular disease or renal disease, such as any of the following (NOT an exhaustive list):

- Hypertension
- Dyslipidemia / Hyperlipidemia
- Diabetes Mellitus
- Congestive Heart Failure (NYHA II-IV)
- Ischemic heart disease
- Peripheral Arterial Disease
- Smoking
- Creatinine Clearance < 60 mL/min or 1 mL/sec
- Acute myocardial infarction, history of myocardial infarction and/or angina
- Stroke, cerebrovascular accident, transient ischemic attacks, and/or amaurosis fugax

If needed, these patients should be treated only after careful consideration (See **WARNINGS AND PRECAUTIONS BOX**).

Endocrine and Metabolism

With nonsteroidal anti-inflammatory treatment there is a potential risk of hyperkalemia, particularly in patients with conditions such as diabetes mellitus or renal failure; elderly patients; or in patients receiving concomitant therapy with beta-adrenergic blockers, angiotensin converting enzyme inhibitors or some diuretics. Serum electrolytes should be monitored periodically during long-term therapy, especially in those patients who are at risk.

Corticosteroids:

GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol) is NOT a substitute for corticosteroids. It does NOT treat corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to exacerbation of corticosteroid-responsive illness. Patients on prolonged corticosteroid therapy should have their therapy tapered slowly if a decision is made to discontinue corticosteroids (see **DRUG INTERACTIONS – Drug-Drug Interactions – Glucocorticoids**).

Gastrointestinal (GI)

The presence of misoprostol in the product may protect against the mucosal damaging effects of the other component, diclofenac.

However, serious GI toxicity, such as peptic/duodenal ulceration, inflammation, perforation, obstruction and gastrointestinal bleeding, sometimes severe and occasionally fatal can occur at any time, with or without symptoms in patients treated with NSAIDs including GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol). NSAIDs, including GD-diclofenac/misoprostol, should be used with caution in patients with a history of, or active, GI

disease, such as ulceration, bleeding, or inflammatory conditions. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and therefore special care should be taken in treating this population. **To minimize the potential risk for an adverse GI event, the lowest effective dose should be used for the shortest possible duration.** For high risk patients, alternate therapies that do not involve NSAIDs should be considered (see **WARNINGS AND PRECAUTIONS – Special Populations – *Geriatrics***).

Minor upper GI problems, such as dyspepsia, commonly occur at any time. Physicians should remain alert for ulceration and bleeding in patients treated with nonsteroidal anti-inflammatory drugs, even in the absence of previous GI tract symptoms.

Physicians should inform patients about the signs and/or symptoms of serious GI toxicity and instruct them to discontinue using GD-diclofenac/misoprostol and seek emergency medical attention if they experience any such symptoms.

Because serious GI tract ulceration and bleeding can occur without warning symptoms, physicians should follow chronically treated patients by checking their hemoglobin periodically and by being vigilant for the signs and symptoms of ulceration and bleeding.

The utility of periodic laboratory monitoring has NOT been demonstrated, nor has it been adequately assessed. Most patients who develop a serious upper GI adverse event on NSAID therapy have no symptoms. Upper GI ulcers, gross bleeding or perforation, caused by NSAIDs, appear to occur in approximately 1% of patients treated for 3-6 months, and in about 2-4% of patients treated for one year. These trends continue, thus increasing the likelihood of developing a serious GI event at some time during the course of therapy. Even short-term therapy has its risks.

Caution should be taken if prescribing GD-diclofenac/misoprostol to patients with a prior history of peptic / duodenal ulcer disease or gastrointestinal bleeding as these individuals have a greater than 10-fold higher risk for developing a GI bleed when taking a NSAID than patients with neither of these risk factors. Other risk factor for GI ulceration and bleeding include the following: *Helicobacter pylori* infection, increased age, prolonged use of NSAID therapy, excess alcohol intake, smoking, female gender, poor general health status or concomitant therapy with any of the following:

- Anti-coagulants (e.g. warfarin)
- Anti-platelet agents (e.g. ASA, clopidogrel)
- Oral corticosteroids (e.g. prednisone)
- Selective Serotonin Reuptake Inhibitors (SSRIs) (e.g. citalopram, fluoxetine, paroxetine, sertraline)

If ulceration is suspected or confirmed, or if GI bleeding occurs, GD-diclofenac/misoprostol should be discontinued immediately, appropriate treatment instituted and the patient monitored closely.

No studies, to date, have identified any group of patients not at risk of developing ulceration and

bleeding. Studies to date show that all NSAIDs can cause GI tract adverse events. Although existing data do not clearly identify differences in risk between various NSAIDs, this may be shown in the future.

Genitourinary

Some NSAIDs are associated with persistent urinary symptoms (bladder pain, dysuria, urinary frequency), hematuria or cystitis. The onset of these symptoms may occur at any time after the initiation of therapy with an NSAID. Some cases have become severe on continued treatment. Should urinary symptoms occur, in the absence of an alternate explanation, treatment with GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol) must be stopped to ascertain if symptoms disappear. This should be done before any urological investigations or treatments are carried out.

Post-menopausal vaginal bleeding may be related to GD-diclofenac/misoprostol administration. If this occurs, diagnostic workup should be undertaken to rule out gynecological pathology (see **ADVERSE DRUG REACTIONS**).

Hematologic

NSAIDs inhibiting prostaglandin biosynthesis interfere with platelet function to varying degrees; patients who may be adversely affected by such an action, such as those on anti-coagulants or suffering from haemophilia or platelet disorders should be carefully monitored when GD-diclofenac/misoprostol is administered.

Anti-coagulants:

The concomitant use of NSAIDs, including diclofenac/misoprostol, with anticoagulants increases the risk of GI and non-GI bleeding and should be given with caution. Concurrent therapy of GD-diclofenac/misoprostol with anticoagulants requires close monitoring of anticoagulation (see **DRUG INTERACTIONS**).

Even with therapeutic INR monitoring, increased bleeding may occur.

Anti-platelet Effects:

NSAIDs inhibit platelet aggregation and have been shown to prolong bleeding time in some patients. Unlike acetylsalicylic acid (ASA), their effect on platelet function is quantitatively less, or of shorter duration, and is reversible. Misoprostol does not exacerbate the effects of diclofenac on platelet activity.

Diclofenac sodium plus misoprostol and other NSAIDs have no proven efficacy as anti-platelet agents and should NOT be used as a substitute for ASA or other anti-platelet agents for prophylaxis of cardiovascular thromboembolic diseases. Anti-platelet therapies (e.g. ASA) should NOT be discontinued. There is some evidence that use of NSAIDs with ASA can markedly attenuate the cardioprotective effects of ASA (see **DRUG INTERACTIONS – Drug-Drug Interactions – Acetylsalicylic Acid (ASA) or other NSAIDs**).

Concomitant administration of GD-diclofenac/misoprostol with low dose ASA increases the risk of GI ulceration and associated complications.

Blood dyscrasias:

Cases of agranulocytosis and hemolytic anemia, some serious, were identified in patients taking diclofenac sodium or diclofenac/misoprostol.

Other blood dyscrasias (such as neutropenia, leucopenia, thrombocytopenia, aplastic anemia) associated with the use of NSAIDs are rare, but could occur with severe consequences.

Anemia is sometimes seen in patients receiving NSAIDs, including GD-diclofenac/misoprostol. This may be due to fluid retention, GI blood loss, or an incompletely described effect upon erythropoiesis. Patients on long-term treatment with NSAIDs, including GD-diclofenac/misoprostol, should have their haemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia or blood loss.

Hepatic/Biliary/Pancreatic

As with other NSAIDs, including GD-diclofenac/misoprostol, borderline elevations of one or more liver function tests (AST, ALT, alkaline phosphatase) may occur in up to 15% of patients. These abnormalities may progress, may remain essentially unchanged, or may be transient with continued therapy. In clinical trials of 4 to 12 weeks duration, clinically significant (>3 times the upper limit of normal) elevations of SGPT (ALT) and/or SGOT (AST), were observed in 2.5% or less of patients who received diclofenac/misoprostol or diclofenac/placebo. In a large trial where patients received diclofenac for a mean of 18 months, ALT/AST elevations >3xULN were observed in 3.1% of patients and elevations >5xULN were observed in 1.3% of patients. ALT/AST elevations usually occur within 1-6 months. However, clinically important liver events, resulting in hospitalization, occurred at various times during the study, and not necessarily early in the course of therapy. Furthermore, more meaningful elevations in transaminases were detected before patients became symptomatic due to routine testing during the trial (see Monitoring and Laboratory testing).

In postmarketing reports of patients receiving diclofenac, cases of drug-induced hepatotoxicity have been reported in the first month, and in some cases, the first 2 months of therapy, but can occur at any time during the treatment. A patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of more severe hepatic reaction while on therapy with this drug. Postmarketing surveillance has reported cases of severe hepatic reactions including jaundice, fulminant hepatitis with and without jaundice, liver necrosis and hepatic failure. Some of these cases have resulted in fatalities or liver transplantation.

Physicians should measure transaminases periodically in patients receiving GD-diclofenac/misoprostol because severe hepatotoxicity may develop without a prodrome of distinguishing symptoms. Severe hepatic reactions can occur at any time during treatment with diclofenac. Although such reactions are rare, if abnormal liver function tests persist or worsen, if clinical signs and symptoms consistent with liver disease develop (e.g. jaundice), or if systemic manifestations occur (e.g. eosinophilia, rash, etc.), this drug should be discontinued immediately.

To minimize the possibility that hepatic injury will become severe between transaminase measurements, physicians should inform patients of the warning signs and symptoms of hepatotoxicity (e.g. nausea, fatigue, lethargy, diarrhea, pruritus, jaundice, right upper quadrant

tenderness, and «flu-like» symptoms), and the appropriate action patients should take if these signs and symptoms appear. Use of GD-diclofenac/misoprostol is contraindicated in patients with significant hepatic impairment or active liver disease. If there is a need to prescribe this drug in the presence of all other patients with liver impairment, it must be done under strict observation.

Caution is advised when using GD-diclofenac/misoprostol in patients with hepatic porphyria, since GD-diclofenac/misoprostol may trigger an attack.

Hypersensitivity Reactions

Anaphylactoid Reactions:

As with NSAIDs in general, anaphylactoid reactions have occurred in patients without known prior exposure to GD-diclofenac/misoprostol. In post-marketing experience, rare cases of anaphylactic/anaphylactoid reactions and angioedema have been reported in patients receiving diclofenac sodium plus misoprostol. GD-diclofenac/misoprostol should NOT be given to patients with the ASA-triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking ASA or other NSAIDs (see **CONTRAINDICATIONS**).

ASA-Intolerance:

GD-diclofenac/misoprostol should NOT be given to patients with complete or partial syndrome of ASA-intolerance (rhinosinusitis, urticaria/angioedema, nasal polyps, asthma) in whom asthma, anaphylaxis, urticaria/angioedema, rhinitis or other allergic manifestations are precipitated by ASA or other NSAIDs. Fatal anaphylactoid reactions have occurred in such individuals. As well, individuals with the above medical problems are at risk of a severe reaction even if they have taken NSAIDs in the past without any adverse reaction (see **CONTRAINDICATIONS**).

Cross-sensitivity:

Patients sensitive to one NSAID may be sensitive to any of the other NSAIDs as well.

Serious skin reactions: (see **WARNINGS AND PRECAUTIONS – Skin**)

Immune

(See **WARNINGS AND PRECAUTIONS – Infection – *Aseptic Meningitis***)

Infection

GD-diclofenac/misoprostol, in common with other NSAIDs, may mask signs and symptoms of an underlying infections disease.

Aseptic Meningitis:

Rarely, with some NSAIDs, the symptoms of aseptic meningitis (stiff neck, severe headaches, nausea and vomiting, fever or clouding of consciousness) have been observed. Patients with autoimmune disorders (systemic lupus erythematosus, mixed connective tissue diseases, etc.) seem to be pre-disposed. Therefore, in such patients, the health care provider must be vigilant to the development of this complication.

Neurologic

Some patients may experience drowsiness, dizziness, blurred vision, vertigo, tinnitus, hearing loss, insomnia or depression with the use of NSAIDs such as GD-diclofenac/misoprostol. If patients experience these side effects, they should exercise caution in carrying out activities that require alertness.

Ophthalmologic

Blurred, diminished vision, and/or sensitivity to light have been reported with the use of NSAIDs. If such symptoms develop, GD-diclofenac/misoprostol should be discontinued and an ophthalmologic examination performed; ophthalmologic examination should be carried out at periodic intervals in any patient receiving GD-diclofenac/misoprostol for an extended period of time.

Peri-Operative Considerations

(See **CONTRAINDICATIONS** – Coronary Artery Bypass Graft Surgery)

Psychiatric

Some patients may experience depression with the use of diclofenac (See **WARNINGS AND PRECAUTIONS** – **Neurologic**).

Renal

Long-term administration of nonsteroidal anti-inflammatory drugs to animals has resulted in renal papillary necrosis and other abnormal renal pathology. In humans, there have been reports of acute interstitial, nephritis, hematuria, low grade proteinuria, and occasionally nephrotic syndrome.

Renal insufficiency due to NSAID use is seen in patients with prerenal conditions leading to the reduction in renal blood flow or blood volume. Under these circumstances, renal prostaglandins help maintain renal perfusion and glomerular filtration rate (GFR). In these patients, administration of a NSAID may cause a reduction in prostaglandin formation and may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with pre-existing renal insufficiency (GFR<60 mL/min or 1 mL/s), dehydrated patients, patients on salt-restricted diets, heart failure, cirrhosis, liver dysfunction, those taking angiotensin-converting enzyme inhibitors, angiotensin-II receptor blockers, cyclosporin, diuretics, and the elderly. Serious or life-threatening renal failure has been reported in patients with normal or impaired renal function after short term therapy with NSAIDs. Even patients at risk who demonstrated the ability to tolerate a NSAID under stable conditions may decompensate during periods of added stress (e.g. dehydration due to gastroenteritis). Discontinuation of NSAIDs is usually followed by recovery to the pre-treatment state.

Diclofenac and its metabolites are eliminated primarily by the kidneys, therefore GD-diclofenac/misoprostol should be used with great caution in patients with impaired renal function. In these cases, utilization of lower doses of diclofenac should be considered and patients carefully monitored. Caution should be used when initiating treatment with NSAIDs, such as GD-diclofenac/misoprostol, in patients with dehydration. It is advisable to rehydrate patients first and then start therapy. The dose should be kept as low as possible and renal function should be monitored.

During long-term therapy kidney function should be monitored periodically.

Advanced Renal Disease: (see **CONTRAINDICATIONS**)

Fluid and Electrolyte Balance:

Use of NSAIDs, such as GD-diclofenac/misoprostol, can promote sodium retention in a dose-dependent manner, which can lead to fluid retention and edema, and consequences of increased blood pressure and exacerbation of congestive heart failure. Thus, caution should be exercised in prescribing GD-diclofenac/misoprostol in patients with a history of congestive heart failure, compromised cardiac function, hypertension, increased age or other conditions predisposing to fluid retention (see **WARNINGS AND PRECAUTIONS – Cardiovascular**).

Use of NSAIDs, such as GD-diclofenac/misoprostol, can increase the risk of hyperkalemia, especially in patients with diabetes mellitus, renal failure, increased age, or those receiving concomitant therapy with adrenergic blockers, angiotensin-converting enzyme inhibitors, angiotensin-II receptor antagonists, cyclosporin, or some diuretics.

Electrolytes should be monitored periodically (see **CONTRAINDICATIONS**).

Respiratory

ASA-induced asthma is an uncommon but very important indication of ASA and NSAID sensitivity. It occurs more frequently in patients with asthma who have nasal polyps.

Sexual Function / Reproduction

The use of GD-diclofenac/misoprostol, as with any drug known to inhibit cyclooxygenase/prostaglandin synthesis, may impair fertility and is not recommended in women attempting to conceive. Therefore, in women who have difficulties conceiving, or who are undergoing investigation of infertility, withdrawal of GD-diclofenac/misoprostol should be considered.

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

Skin

In rare cases, serious skin reactions such as Stevens-Johnson syndrome, toxic epidermal necrolysis, exfoliative dermatitis and erythema multiforme have been associated with the use of some NSAIDs. Because the rate of these reactions is low, they have usually been noted during post-marketing surveillance in patients taking other medications also associated with the potential development of these serious skin reactions. Thus, causality is NOT clear. These reactions are potentially life-threatening but may be reversible if the causative agent is discontinued and appropriate treatment instituted. Patients should be advised that if they experience a skin rash they should discontinue their NSAID and contact their physician for assessment and advice, including which additional therapies to discontinue.

Serious skin reactions, some of them fatal, including drug reaction with eosinophilia and systemic symptoms (DRESS syndrome), exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs, including GD-diclofenac/misoprostol (see **ADVERSE REACTIONS - Post-Market Adverse Drug Reactions**). Patients appear to be at higher risk for these events early in the course of therapy: the onset of the event occurring in the majority of cases within the first month of treatment. GD-diclofenac/misoprostol should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

GD-diclofenac/misoprostol may cause sensitivity to sunlight. Any exposure to sunlight or sunlamps may cause sunburn, skin blisters, skin rash, redness, itching or discoloration. Patients should be advised that if they experience any of these symptoms, they should discontinue their NSAID and contact their physician for assessment and advice, including which additional therapies to discontinue.

Special Populations

Pregnant Women:

GD-diclofenac/misoprostol is CONTRAINDICATED for use in women who are **pregnant**, or in whom **pregnancy has not been excluded**. Misoprostol administration to pregnant women induces uterine contractions and is associated with abortion, premature birth, birth defects and fetal death. Misoprostol can cause uterine tetany and uterine rupture if administered to pregnant women beyond the eighth week of pregnancy (see **CONTRAINDICATIONS and ADVERSE REACTIONS, – Post-Market Adverse Drug Reactions**).

Inhibition of prostaglandin synthesis may adversely affect pregnancy and/or the embryo-foetal development. Data from epidemiological studies suggest an increased risk of miscarriage and of cardiac malformation after use of a prostaglandin synthesis inhibitor in early pregnancy.

NSAIDs including diclofenac may result in reduction of amniotic fluid volume and even oligohydramnios. Such effects may occur within a few days after treatment initiation and are usually reversible. The rate of oligohydramnios after treatment with some NSAIDs for 2 to 8 weeks was reported as high as 38% or even higher. NSAIDs were also shown to cause significant reduction in fetal urine production prior to reduction of amniotic fluid volume.

In animals, administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre-and post-implantation loss and embryo-foetal lethality. In addition, increased incidences of various malformations, including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period.

Nursing Women:

Misoprostol is rapidly metabolised in the mother to misoprostol acid, which is biologically active and is excreted in breast milk. Diclofenac/misoprostol is contraindicated in nursing mothers because the excretion of misoprostol acid could cause undesirable effects such as diarrhea in nursing infants (see **CONTRAINDICATIONS**).

Pediatrics (<18 years of age):

(See **CONTRAINDICATIONS**)

Geriatrics (>65 years of age):

Patients older than 65 years and frail or debilitated patients are most susceptible to a variety of adverse reactions from NSAIDs. The incidence of these adverse reactions increases with dose and duration of treatment. In addition, these patients are less tolerant to ulceration and bleeding. Most reports of fatal GI events are in this population. Older patients are also at risk of lower esophageal ulceration and bleeding (see **ACTION AND CLINICAL PHARMACOLOGY**).

For such patients, consideration should be given to a starting dose lower than the one usually recommended, with individual adjustment when necessary and under close supervision.

As with any NSAID, the elderly are likely to tolerate adverse events less well than younger patients.

Diclofenac is known to be substantially excreted by the kidney, and the risk of toxic reactions to GD-diclofenac/misoprostol may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function (see **WARNINGS AND PRECAUTIONS – Renal**).

Monitoring and Laboratory Tests

Cardiovascular: (Hypertension): Blood pressure should be monitored regularly during therapy with GD-diclofenac/misoprostol.

Hematologic: Patients on long-term treatment with NSAIDs, including GD-diclofenac/misoprostol, should have their hemoglobin, hematocrit, red blood cells, white blood cells, and platelets checked if they exhibit any signs or symptoms of anemia or blood loss or blood dyscrasia.

Concurrent therapy of GD-diclofenac/misoprostol with warfarin requires close monitoring of the international normalized ratio (INR).

Hepatic: Hepatic functions (e.g. serum transaminases, bilirubine) should be performed within 4 to 8 weeks of starting therapy, and then monitored regularly during therapy with GD-diclofenac/misoprostol. Patients with symptoms and/or signs of liver dysfunction, or in whom an

abnormal liver function test has occurred, their hepatic function (e.g. serum transaminases, bilirubin) should be monitored carefully for evidence of the development of a more severe hepatic reaction while on therapy with GD-diclofenac/misoprostol. If abnormal liver tests persist or worsen, GD-diclofenac/misoprostol should be discontinued.

Ophthalmologic: Patients on long-term treatment with **GD-diclofenac/misoprostol** should have an ophthalmologic examination performed periodically, and if they experience blurred and/or diminished vision.

Renal: Renal function should be monitored in high-risk populations, such as the elderly, patients with advanced renal disease, patients with cardiovascular disease and diabetes mellitus, as well as in the setting of concomitant use of diuretics and ACE inhibitors (see **CONTRAINDICATIONS**). If abnormal renal tests persist or worsen, GD-diclofenac/misoprostol should be discontinued.

Electrolytes, including serum potassium, should be monitored periodically, especially in patients with diabetes mellitus, renal failure, increased age, or those receiving concomitant therapy with adrenergic blockers, angiotensin-converting enzyme inhibitors, angiotensin-II receptor antagonists, cyclosporine, or some diuretics.

Laboratory abnormalities included increased alkaline phosphatase, decreased hematocrit and elevated SGPT (ALT).

Persistently abnormal or worsening renal, hepatic or hematological test values should be followed up carefully since they may be related to therapy.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

The most common adverse reactions encountered with nonsteroidal anti-inflammatory drugs are gastrointestinal, of which peptic ulcer, with or without bleeding, is the most severe. Fatalities have occurred, particularly in the elderly.

Most fatal gastrointestinal events occur in the elderly or debilitated patients. Gastrointestinal adverse events can develop at any time in the course of the therapy.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

In clinical trials, 3549 arthritic patients have been treated with diclofenac sodium plus misoprostol, 506 of whom received diclofenac sodium plus misoprostol for more than one year. A total of 285 patients have been treated with diclofenac sodium plus misoprostol 75 in clinical trials for a duration of up to 12 weeks.

The following adverse reactions occurred with an incidence of 1% or greater with at least one of the diclofenac sodium plus misoprostol or Diclofenac dosing regimens presented below:

	A50 ¹ BID N=391	A50 ¹ TID N=692	A50 ¹ BID/TID N=750	D50 ² BID/TID N=754	A75 ³ BID* N=285	D75 ⁴ BID* N=260
Body as a Whole:						
Influenza-Like Symptoms	1.0	0.6	2.0	1.5	1.1	2.3
Pain	0.5	1.0	0.7	0.8	4.2	1.9
Back Pain	0.8	0.6	1.2	1.1	3.2	3.5
Chest Pain	1.0	0.3	1.1	0.5	0.7	3.1
Fever	0.0	0.6	0.7	1.1	1.4	0.0
Asthenia	0.0	0.1	0.1	0.7	1.1	0.4
Myalgia	0.8	0.7	0.8	0.3	1.1	0.4
Arthralgia	0.0	0.7	0.3	0.3	2.8	3.5
Arthrosis	0.0	0.1	0.3	0.0	1.4	1.2
Cardiovascular:						
Hypertension	0.0	0.5	0.0	0.1	1.1	2.3
Edema	0.8	0.7	0.0	0.3	1.1	1.2
Dependent Edema	0.0	0.3	0.4	0.5	1.1	0.4
Leg Edema	0.0	0.1	0.0	0.1	1.1	0.8
Central Nervous System:						
Headache	9.2	6.4	7.3	9.2	12.3	15.8
Dizziness	2.6	2.0	3.5	5.3	3.9	4.2
Migraine	1.3	0.6	0.4	0.9	1.4	0.8
Paresthesia	0.3	0.3	0.7	0.7	1.1	0.4
Dermatologic:						
Rash	0.8	1.4	1.5	1.1	2.1	3.5
Pruritis	1.0	0.4	1.2	0.9	2.1	1.9
Skin ulceration	0.0	0.0	0.1	0.0	1.1	0.4
Gastrointestinal:						
Abdominal Pain	19.4	19.4	23.2	19.5	24.6	24.2
Diarrhea	15.9	17.8	19.9	11.3	20.4	16.2
Dyspepsia	7.2	14.5	11.3	7.8	33.3	34.6
Nausea	10.2	10.0	11.7	6.5	14.0	9.2
Flatulence	6.1	8.7	8.0	3.1	18.2	9.2
Gastritis	2.8	2.3	3.6	6.8	7.4	13.1
Vomiting	2.6	3.3	3.1	1.3	3.9	5.4
Constipation	1.8	2.6	2.1	2.9	4.9	6.9
Eructation	2.6	0.3	2.0	0.8	2.1	0.4

	A50¹ BID N=391	A50¹ TID N=692	A50¹ BID/TID N=750	D50² BID/TID N=754	A75³ BID* N=285	D75⁴ BID* N=260
Esophagitis	0.8	1.7	1.1	0.8	3.9	1.9
Duodenitis	2.3	0.9	0.9	2.3	3.5	5.0
Gastroesophageal Reflux	0.0	1.0	0.4	1.7	1.1	1.2
Duodenal Ulcer	0.0	1.2	0.1	0.4	0.7	2.7
Gastric Ulcer	0.8	0.6	0.7	1.7	3.2	6.9
Tooth Disorder	0.3	0.6	0.0	0.0	1.1	0.8
Hepatic:						
SGPT (ALT) Increased	0.5	0.6	0.1	0.7	2.5	2.3
SGOT (AST) Increased	0.5	0.4	0.0	0.5	1.1	2.3
Psychiatric:						
Insomnia	1.3	0.4	0.9	1.2	2.5	1.9
Somnolence	0.8	0.6	0.7	0.9	1.1	0.8
Respiratory:						
Upper Respiratory Tract Infection	1.0	2.7	1.1	2.1	2.8	3.8
Pharyngitis	0.5	1.9	1.1	1.9	3.5	1.5
Rhinitis	0.8	2.6	0.3	0.9	3.2	4.2
Sinusitis	0.0	0.9	0.1	0.1	6.0	2.7
Coughing	0.3	1.2	0.4	1.2	1.8	3.5
Bronchitis	0.0	0.4	0.7	1.1	2.1	1.5
Dyspnea	0.3	0.4	0.4	0.7	1.4	0.4
Urogenital:						
Menorrhagia	0.9	0.6	1.3	0.0	0.0	0.5
Vaginitis	0.0	0.9	0.0	0.0	1.0	1.1
Perineal Pain, male	0.6	0.0	0.0	0.0	1.1	0.0

¹ A50 = diclofenac sodium plus misoprostol 50

² D50 = Diclofenac 50mg

³ A75 = diclofenac sodium plus misoprostol 75

⁴ D75 = Diclofenac 75 mg

*Patients must have experienced ulceration in order to enter study. This represents an extremely high risk cohort

Abdominal pain and diarrhea were generally transient and mild to moderate in severity, occurring early in the course of therapy and lasting several days. The abdominal pain and diarrhea usually resolved spontaneously while continuing diclofenac sodium plus misoprostol.

Less Common Clinical Trial Adverse Drug Reactions (<1%)

The following adverse events were reported by 1% or less of the subjects receiving diclofenac sodium plus misoprostol. Causal relationships between diclofenac sodium plus misoprostol and these events have not been established but cannot be excluded.

Body as a Whole:	hot flushes, malaise, rigors
Cardiovascular:	palpitation and syncope
Central Nervous System/ Psychiatric:	anorexia, anxiety, concentration impaired, depression, hypoesthesia, speech disorder and vertigo
Dermatologic:	angioedema, erythematous rash, sweating increased, urticaria and purpura
Gastrointestinal:	mouth dry, abdomen enlarged, esophageal ulceration, glossitis, hematemesis, hiccup and melena
Gynecological:	menstrual disorder, intermenstrual bleeding, dysmenorrhea, leukorrhea, vaginal bleeding, breast pain and uterine cramping. (Post-menopausal vaginal bleeding may be related to GD-diclofenac/misoprostol administration. If this occurs, diagnostic workup should be undertaken to rule out gynecological pathology.)
Hematologic:	leukopenia and thrombocytopenia
Hepatic:	gall bladder disorder, bilirubinemia, abnormal hepatic function, LDH increased, and alkaline phosphatase increased, hepatitis
Metabolic:	BUN increased and glycosuria
Respiratory:	hyperventilation and sputum increased
Special Senses:	earache, eye pain, taste loss, taste abnormalities, tinnitus and vision abnormal
Urinary:	Dysuria and urine abnormal

Abnormal Hematologic and Clinical Chemistry findings

See **DRUG INTERACTIONS - Drug Laboratory Interactions**

Post-Market Adverse Drug Reactions

Additional reports of serious adverse events temporally associated with diclofenac sodium plus misoprostol during worldwide post-marketing experience are included below. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to diclofenac sodium plus misoprostol exposure.

Body as a whole:	Death, fatigue, infection, sepsis
Immune System Disorders:	Allergic reactions including anaphylaxis and angioedema, Laryngeal/pharyngeal edema

Cardiovascular:	Myocardial infarction, stroke, transient ischemic attack, cerebral hemorrhage, hypertension, cardiac failure, vasculitis arrhythmia, atrial fibrillation, congestive heart failure, hypotension, increased CPK, phlebitis, premature ventricular contractions, tachycardia
Central Nervous System/ Psychiatric:	Changes in mood, nightmares, meningitis aseptic, tremor, coma, convulsions, drowsiness, hyperesthesia, hypertonia, neuralgia, confusion, disorientation, dream abnormalities, hallucinations, irritability, nervousness, paranoia, psychotic reaction
Dermatologic:	Cutaneous reactions (including rash, pruritus and bullous eruption), rare cases of mucocutaneous reactions, Edema, urticaria, Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme, dermatitis exfoliative acne, alopecia, bruising, pemphigoid reaction, photosensitivity, pruritus ani, drug reaction with eosinophilia and systemic symptoms (DRESS syndrome)
Gastrointestinal:	Pancreatitis, stomatitis and ulcerative stomatitis, gastrointestinal inflammation, gastrointestinal bleeding, gastrointestinal ulceration, gastrointestinal perforation, gastrointestinal neoplasm benign, heartburn, hemorrhoids, tenesmus, appetite changes, dry mouth, dysphagia, enteritis
Gynecological:	Abnormal uterine contractions, uterine hemorrhage, uterine rupture/perforation, retained placenta, amniotic fluid embolism, incomplete abortion, premature birth, fetal death, and birth defects, female fertility decreased, reduction of amniotic fluid volume, reduction of fetal urine production
Hematology:	Thrombocytopenia, platelet aggregation inhibition, hemolytic anemia, agranulocytosis, anemia, aplastic anemia, coagulation time increased, ecchymosis, eosinophilia, epistaxis, leukocytosis, lymphadenopathy, pancytopenia, pulmonary embolism, rectal bleeding, thrombocythemia, decreased hematocrit
Hepatic:	Hepatitis, hepatotoxicity, severe hepatic reactions including liver necrosis, jaundice, fulminant hepatitis with and without jaundice, and liver failure, with a fatal outcome or requiring liver transplantation (see WARNINGS AND PRECAUTIONS – Hepatic/Biliary/Pancreatic).
Special Senses:	Blurred vision, hearing impairment, amblyopia, conjunctivitis, diplopia, glaucoma, iritis, lacrimation abnormal, night blindness

Urogenital:	Renal failure, interstitial nephritis, glomerulonephritis, glomerulonephritis membranous, glomerulonephritis minimal lesion, renal papillary necrosis, nephrotic syndrome, renal impairment, impotence, cystitis, hematuria, micturition frequency, nocturia, oliguria/polyuria, proteinuria, urinary tract infection
Metabolism and nutrition disorders:	Fluid retention, alkaline phosphatase increased, dehydration, gout, hypercholesterolemia, hyperglycemia, hyperuricemia, hypoglycemia, hyponatremia, periorbital edema, porphyria, weight changes
Respiratory system:	Asthma, pneumonia, respiratory depression

Meta-analysis and pharmacoepidemiological data point towards an increased risk of arteriothrombotic events associated with the use of diclofenac, particularly at a high dose (see **WARNINGS AND PRECAUTIONS BOX**).

DRUG INTERACTIONS

Overview

Factors such as excess alcohol intake, smoking, and concomitant NSAID and oral steroid or anticoagulant use have been associated with increased risk of GI adverse events such as ulceration and bleeding. In laboratory studies, misoprostol has shown no significant effect on the cytochrome P450-linked hepatic mixed function oxidase system, and therefore should not affect the metabolism of theophylline, warfarin, benzodiazepines or other drugs normally metabolized by this system.

Diclofenac metabolism is predominantly mediated via cytochrome P450 CYP 2C9 in the liver. Patients who are known or suspected to be poor CYP2C9 metabolizers based on previous history/experience with other CYP2C9 substrates should be administered diclofenac with caution as they may have abnormally high plasma levels due to reduced metabolic clearance. Caution is recommended when co-prescribing diclofenac with potent CYP2C9 inhibitors (such as sulfapyrazole and voriconazole), which could result in a significant increase in peak plasma concentrations and exposure to diclofenac due to inhibition of diclofenac metabolism. Caution should be exercised when prescribing GD-diclofenac/misoprostol with concomitant drugs that are known to be potentially hepatotoxic (e.g. antibiotics, anti-epileptics).

Drug-Drug Interactions

Misoprostol has been used concomitantly with at least 44 different classes of drugs, including more than 150 drugs. There were no reports of any clinically significant drug interactions.

Acetylsalicylic Acid (ASA) or other NSAID's:

When diclofenac and ASA are taken simultaneously, the bioavailability of each is reduced. Concomitant administration of GD-diclofenac/misoprostol and ASA is not recommended because diclofenac is displaced from its binding sites by ASA, resulting in lower plasma concentrations, peak plasma levels and AUC values. Misoprostol does not affect the kinetics of other NSAIDs (e.g., ibuprofen, indomethacin and piroxicam). The use of GD-diclofenac/misoprostol in

addition to any other NSAID, including those over the counter ones (such as ASA and ibuprofen) for analgesic and/or anti-inflammatory effects is NOT recommended because of the absence of any evidence demonstrating synergistic benefits and the potential for additive adverse reactions.

The exception is the use of low dose ASA for cardiovascular protection, when another NSAID is being used for its analgesic/anti-inflammatory effect, keeping in mind that combination NSAID therapy is associated with additive adverse reactions.

Some NSAIDs (e.g. ibuprofen) may interfere with the anti-platelet effects of low dose ASA, possibly by competing with ASA for access to the active site of cyclooxygenase-I.

Antacids:

Only aluminum-based antacids should be used with GD-diclofenac/misoprostol as magnesium-based antacids may increase the potential for diarrhea (see **ADVERSE REACTIONS**). The concomitant administration of aluminum hydroxide or magnesium hydroxide antacids may delay the absorption of diclofenac but does not affect the total amount of the drug absorbed. The total availability of misoprostol acid is reduced by antacids in large doses.

Anticoagulants:

Numerous studies have shown that the concomitant use of NSAIDs and anticoagulants increases the risk of GI adverse events such as ulceration and bleeding. Pharmacodynamic studies have shown no potentiation of anticoagulant drugs due to concurrent administration with diclofenac. However, other NSAIDs have been shown to interact with anticoagulant agents. Although clinical investigations would appear to indicate that diclofenac has no influence on the effect of anticoagulants, there are isolated reports of an increased risk of hemorrhage with the combined use of diclofenac and nicoumalone anticoagulant therapy. Special caution is therefore recommended and frequent laboratory tests should be performed to check that the desired response to the anticoagulant is being maintained. Because prostaglandins play an important role in hemostasis and NSAIDs affect platelet function as well, concurrent therapy of GD-diclofenac/misoprostol with warfarin requires close monitoring to be certain no change in anticoagulant dosage is necessary (see **WARNINGS AND PRECAUTIONS – Hematologic – Anti-coagulants**).

Anti-hypertensives:

NSAIDs may diminish the anti-hypertensive effect of diuretics and other antihypertensive drugs including Angiotensin Converting Enzyme (ACE) inhibitors, angiotensin II antagonists (AIIA) and beta-blockers.

Co-administration of ACE inhibitors, angiotensin-II antagonists, or diuretics with NSAIDs can have an increased risk for deterioration of renal function, acute renal failure and hyperkalemia. Blood pressure and renal function (including electrolytes) should be monitored more closely in this situation, as there can be a substantial increase in blood pressure.

Dehydrated patients or elderly patients with compromised renal function may be at greater risk.

Anti-platelet Agents (including ASA):

There is an increased risk of bleeding, via inhibition of platelet function, when anti-platelet agents are combined with NSAIDs, such as GD-diclofenac/misoprostol (see **WARNINGS AND**

PRECAUTIONS – Hematologic – *Anti-platelet Effects*).

Cyclosporin or Tacrolimus:

When co-administered with cyclosporine, there is a two-fold increase in diclofenac systemic exposure. It is prudent to start with the lowest dose of diclofenac/misoprostol and to monitor closely for signs of toxicity.

Co-administration of cyclosporin or tacrolimus may also increase the nephrotoxic effect of cyclosporin or tacrolimus due to the NSAID's effect on renal prostaglandins. Renal function should be monitored when GD-diclofenac/misoprostol and either of these drugs is used in combination.

Digoxin:

Diclofenac may increase the plasma concentration of digoxin. Dosage adjustment of the digoxin may be required with GD-diclofenac/misoprostol. Serum digoxin levels should be monitored for possible digoxin toxicity.

Diuretics/Antihypertensives:

Clinical studies as well as post-marketing observations have shown that NSAIDs can reduce the effect of diuretics. Concomitant treatment of GD-diclofenac/misoprostol with potassium-sparing diuretics may be associated with increased serum potassium levels, thus making it necessary to monitor the latter. The antihypertensive effect of hydrochlorothiazide and ACE inhibitors may be decreased by diclofenac in patients with essential hypertension. Co-administration of GD-diclofenac/misoprostol with ACE inhibitors may result in an impairment of renal function.

Glucocorticoids:

Some studies have shown that the concomitant use of NSAIDs and oral glucocorticoids increases the risk of GI side effects such as ulceration and bleeding. This is especially the case in older (>65 years of age) individuals.

Lithium:

Diclofenac, when administered concomitantly with lithium, increases the lithium plasma concentration through an effect on lithium renal clearance. Lithium toxicity may develop in these patients. Dosage adjustment of lithium may be required with GD-diclofenac/misoprostol.

Methotrexate:

Concurrent administration of methotrexate and diclofenac may result in increased plasma levels of methotrexate and rare cases of fatal renal toxicity have been reported. Thus, caution should be taken when administering GD-diclofenac/misoprostol and methotrexate.

Oral Contraceptives:

No drug interaction data are available for ARTHROTEC and the co-administration of oral contraceptives.

Oral hypoglycemic agents:

Diclofenac does not alter glucose metabolism in normal subjects, and pharmacodynamic studies have shown no potentiation of oral hypoglycemic drugs due to concurrent administration with diclofenac. However, other NSAIDs have been shown to interact with oral hypoglycemic

agents. Therefore, GD-diclofenac/misoprostol should be administered with caution in patients receiving insulin or oral hypoglycemic agents.

Phenytoin:

When using phenytoin concomitantly with diclofenac, monitoring of phenytoin plasma concentrations is recommended due to an expected increase in exposure to phenytoin.

Selective Serotonin Reuptake Inhibitors (SSRIs):

Concomitant administration of NSAIDs and SSRIs may increase the risk of gastrointestinal ulceration and bleeding (see **WARNINGS AND PRECAUTIONS – Gastrointestinal**).

Sulfinpyrazone:

Concomitant administration of diclofenac and sulfinpyrazone could result in a significant increase in peak plasma concentrations and exposure to diclofenac due to inhibition of diclofenac metabolism.

Voriconazole:

Voriconazole increased C_{max} and AUC of diclofenac (50 mg single dose) by 114% and 78%, respectively.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Diclofenac increases platelet aggregation time but does not affect bleeding time, plasma prothrombin clotting time, plasma fibrinogens, or factors V and VII to XII. Statistically significant changes in prothrombin and partial thromboplastin times have been reported in normal volunteers. The mean changes were observed to be less than 1 second in both instances, and are unlikely to be clinically important.

Drug-Lifestyle Interactions

Smoking and alcohol intake should be discouraged while taking GD-diclofenac/misoprostol as they constitute risk factors for increased cardiovascular and gastrointestinal problems respectively. Patients experiencing visual disturbances, dizziness, vertigo, somnolence or other central nervous system disturbances while taking **GD-diclofenac/misoprostol** should refrain from driving or using machines.

DOSAGE AND ADMINISTRATION

Dosing Considerations

In elderly patients: the dosage should be reduced to the lowest dose that will provide control of symptoms, adjusted when necessary, and closely supervised (see **WARNINGS AND PRECAUTIONS – Special Populations – Geriatrics**).

Cardiovascular disease or cardiovascular risk factors: Treatment with GD-diclofenac/misoprostol is not recommended in patients with pre-existing cardiovascular disease (congestive heart failure NYHA II-IV, ischemic heart disease, peripheral arterial disease), cerebrovascular disease, uncontrolled hypertension, or patients with risk factors for cardiovascular disease (e.g. hypertension, hyperlipidemia, diabetes mellitus and smoking). These patients should be treated with GD-diclofenac/misoprostol only after careful consideration (see **WARNINGS AND PRECAUTIONS-BOX**).

Renal Insufficiency: In patients with mild to moderate renal insufficiency, the lowest dose of GD-diclofenac/misoprostol should be considered, and patients should be monitored closely (see **WARNINGS AND PRECAUTIONS – Renal**). GD-diclofenac/misoprostol is contraindicated in patients with severe renal impairment (estimated creatinine clearance < 30 mL/min or 0.5mL/sec) (see **CONTRAINDICATIONS**).

Hepatic Insufficiency: If GD-diclofenac/misoprostol must be used in patients with mild to moderate hepatic impairment, these patients must be closely monitored (see **WARNINGS AND PRECAUTIONS – Hepatic/Biliary/Pancreatic**). GD-diclofenac/misoprostol is contraindicated in patients with significant hepatic impairment or active liver disease (see **CONTRAINDICATIONS**). Diclofenac metabolism is predominantly mediated via cytochrome P450 CYP 2C9 in the liver. Patients who are known or suspected to be poor CYP2C9 metabolizers based on previous history/experience with other CYP2C9 substrates should be administered diclofenac with caution (see **DRUG INTERACTIONS – Overview**). Caution should be exercised when prescribing GD-diclofenac/misoprostol with concomitant drugs that are known to be potentially hepatotoxic (e.g. antibiotics, anti-epileptics).

Recommended Dose and Dosage Adjustment

Use of GD-diclofenac/misoprostol should be limited to the lowest effective dose in every patient (see **WARNINGS AND PRECAUTIONS**).

The recommended daily oral dose of GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol) for treating the signs and symptoms of rheumatoid arthritis and osteoarthritis is 100 mg administered as two divided doses (50 mg twice per day) (see **WARNINGS AND PRECAUTIONS – Cardiovascular**).

The recommended maximum daily dose is 100 mg.

GD-diclofenac/misoprostol should be taken ***immediately after a meal or with food or milk*** and the tablets should be swallowed whole.

Missed Dose

If a dose of GD-diclofenac/misoprostol is missed, the next dose should be taken at the regular time. The dose should not be doubled.

OVERDOSAGE

Diclofenac Sodium

Worldwide reports on overdosage with diclofenac cover 27 cases. In 10 of these 27 cases, diclofenac was the only drug taken; all of these patients recovered. The highest dose of diclofenac was 2.5 g in a 20-year-old male who suffered acute renal failure as a consequence, and who was treated with dialysis sessions and recovered in 2 days. The next highest dose was 2.35 g in a 17-year-old female who experienced vomiting and drowsiness. A dose of 2.0 g of diclofenac was taken by a woman of unspecified age who remained asymptomatic.

There is no specific antidote for diclofenac. In cases of overdosage, absorption should be prevented as soon as possible by means of induction of vomiting, gastric lavage or treatment with activated charcoal.

Supportive and symptomatic treatment should be given for complications such as drowsiness, confusion, general hypotonia, hypotension, renal failure, convulsions, gastrointestinal irritation and respiratory depression. Measures to accelerate elimination (forced diuresis, hemoperfusion, dialysis) may be considered, but may be of limited use because of the high protein-binding and extensive metabolism (diclofenac 99% protein bound and misoprostol acid less than 90% protein bound).

Misoprostol

The toxic dose of misoprostol in humans has not been determined. Cumulative total daily doses of 1,600 µg have been tolerated with only symptoms of gastrointestinal discomfort being reported.

Clinical signs that may indicate an overdose are sedation, tremor, convulsions, dyspnea, abdominal pain, diarrhea, fever, palpitations, hypotension, or bradycardia. Symptoms should be treated with supportive therapy.

It is not known if misoprostol acid is dialyzable. However, because misoprostol is metabolized like a fatty acid, it is unlikely that dialysis would be appropriate treatment for overdosage.

The use of oral activated charcoal may help to reduce the absorption of diclofenac and misoprostol.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol) is a combination of a

nonsteroidal anti-inflammatory drug (NSAID) with a mucosal protective synthetic analog of prostaglandin E₁.

Diclofenac inhibits prostaglandin synthesis by interfering with the action of prostaglandin synthetase. This inhibitory effect may partially explain its actions, both therapeutic and adverse.

Misoprostol has been shown to inhibit both basal and stimulated gastric acid secretion. In addition, increases in gastric mucosal blood flow, duodenal bicarbonate secretion and gastric mucus secretion have all been observed following treatment with misoprostol. It is not known whether the ability of misoprostol to prevent gastric and duodenal ulcers is the result of its antisecretory effect, its mucosal protective effect, or both.

Pharmacokinetics

The pharmacokinetic profiles of diclofenac and misoprostol administered alone are similar to the profiles when they are coadministered as separate tablets, or given as GD-diclofenac/misoprostol (diclofenac sodium plus misoprostol). No pharmacokinetic interaction between the two drugs has been observed following either single or multiple doses.

There was no accumulation of diclofenac or misoprostol acid in plasma following repeated doses of diclofenac sodium plus misoprostol.

Table 1 Summary of Mean Diclofenac Sodium plus Misoprostol Pharmacokinetic Parameters

	C _{max}		AUC		T _{max}	
	D* (µg/mL)	M* (pg/mL)	D (µg.h/mL)	M (pg.h/mL)	D (h)	M (h)
Single dose of diclofenac sodium plus misoprostol 50, Healthy Male Subjects, N=36	1.13	136	1.63 (AUC ₍₀₋₂₄₎)	238 (AUC ₍₀₋₄₎)	3.9	0.87
Single dose of diclofenac sodium plus misoprostol 75, Healthy Male and Female Subjects, N=35	2.03	304	2.77 (AUC ₍₀₋₁₂₎)	177 (AUC ₍₀₋₄₎)	1.96	0.26

* D = Diclofenac; M = Misoprostol acid

Absorption:

Orally administered diclofenac is rapidly and almost completely absorbed.

Orally administered misoprostol is also rapidly and extensively absorbed.

With diclofenac sodium plus misoprostol the effect of food on the bioavailability of the diclofenac and misoprostol components is similar to that reported for the individual drugs. The times of peak concentration (T_{max}) for diclofenac and misoprostol are prolonged by approximately 50% and 100% respectively, while the peak concentrations (C_{max}) are decreased by about 25% for diclofenac and 50% for misoprostol: the AUC for diclofenac is decreased by approximately 60%, while that of misoprostol is increased by about 25%.

Distribution:

Diclofenac is highly but reversibly bound in the plasma. Following administration of enteric-

coated tablets there is high between- and within-subject variability in the plasma concentrations of diclofenac, particularly if the tablets are taken with food. However, the plasma concentrations show a linear relationship to the amount of drug administered and no accumulation occurs provided that the recommended dosage intervals are observed.

There is high variability in plasma levels of misoprostol acid, but mean values after single doses show a linear relationship with dose over the range of 200 to 400 µg. No accumulation has been found in multiple dose studies and plasma steady state was achieved within two days.

Metabolism:

Diclofenac metabolism is predominantly mediated via cytochrome P450 CYP 2C9 in the liver. Diclofenac undergoes single and multiple hydroxylation and methoxylation, producing 3'-, 4'-, 5-hydroxy, 4'-5-hydroxy and 3'-hydroxy-4'-methoxy derivatives of diclofenac. These phenolic metabolites are largely inactive, and (along with the parent compound) are mostly converted to glucuronide conjugates.

Misoprostol undergoes rapid metabolism to misoprostol acid.

Excretion:

The half-life of diclofenac is 1 to 2 hours. Forty to 60% of the drug and its metabolites are eliminated in the urine and the balance in the bile.

Misoprostol acid is quickly eliminated (elimination half-life of approximately 30 minutes). Approximately 70% of the dose of misoprostol is excreted in the urine, mainly as biologically inactive metabolites.

Special Populations and Conditions

Geriatrics:

The kinetics and metabolism of diclofenac do not appear to be affected by age. In the elderly, the AUC of misoprostol acid is increased by roughly 40%.

Poor CYP2C9 Metabolizers:

Diclofenac metabolism is predominantly mediated via cytochrome P450 CYP 2C9 in the liver.

Patients who are known or suspected to be poor CYP2C9 metabolizers based on previous history/experience with other CYP2C9 substrates should be administered diclofenac with caution (see **DRUG INTERACTIONS – Overview**).

Hepatic Insufficiency:

The kinetics and metabolism of diclofenac do not appear to be affected by hepatic impairment.

Misoprostol does not affect the hepatic mixed function oxidase (cytochrome P-450) enzyme system in animals. In a study of people with mild to moderate hepatic impairment, mean misoprostol acid AUC and C_{max} showed approximately double the mean values obtained in healthy people. Three people who had the lowest antipyrine and lowest indocyanine green clearance values had the highest misoprostol acid AUC and C_{max} values.

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

Renal Insufficiency:

Differences in the pharmacokinetics of diclofenac (50 mg intravenously) have not been detected in studies of patients with renal impairment (N=5, creatinine clearance 3 to 42 mL/min). In these patients, AUC values and elimination rates were comparable to those in healthy people.

Following oral administration of misoprostol in patients with mild-to-moderate renal impairment, there was no significant effect on the pharmacokinetic profile compared to normal subjects. However, in anuric patients, an approximate doubling of C_{max} , AUC and $t_{1/2}$ of misoprostol acid has been observed compared to normal subjects.

STORAGE AND STABILITY

Store at 15 to 25°C and protect from heat and humidity.

DOSAGE FORMS, COMPOSITION AND PACKAGING

GD-diclofenac/misoprostol 50 tablets (diclofenac sodium plus misoprostol) are white to off-white, round biconvex tablets, engraved "SEARLE" over "1411" on one side, 4 x "A" around the circumference of the reverse side with a "50" in the middle. Each tablet has an enteric-coated core containing 50 mg diclofenac sodium, surrounded by an outer mantle containing 200 µg misoprostol.

Bottles of 250.

GD-diclofenac/misoprostol 75 tablets are white to off-white, round and biconvex, engraved "SEARLE" over "1421" on one side, 4 x "A" around the circumference of the reverse side with a "75" in the middle. Each tablet has an enteric-coated core containing 75 mg diclofenac sodium, surrounded by an outer mantle containing 200 µg misoprostol.

Bottles of 250.

Non-medicinal Ingredients:

GD-diclofenac/misoprostol 50 and GD-diclofenac/misoprostol 75 contain: Hydrogenated Castor Oil, Microcrystalline Cellulose, Colloidal Silicon Dioxide, Corn Starch, Crospovidone, Hypromellose, Lactose, Magnesium Stearate, Methacrylic Acid Copolymer, Povidone K-30, Sodium Hydroxide, Talc, Triethyl Citrate.

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Proper name: Diclofenac Sodium

Chemical name: Sodium [*o*-(2,6-dichloroanilino) phenyl acetate]

Molecular formula: C₁₄H₁₀Cl₂NO₂Na

Molecular weight: 318.1

Structural formula:



Physicochemical

properties:

Diclofenac sodium is a white to off-white powder with a salty, bitter taste. At 25°C, diclofenac sodium is 2% soluble in water (pH 7.7). It is practically insoluble in aqueous acidic solutions.

pH: 7.2

Melting Point: 280°-290°C with decomposition

pKa: 3.8 (potentiometry)
4.7 (spectrophotometry)

Drug Substance


Proper name: Misoprostol

Chemical name: (±)-(11 α , 13E)-11,16-dihydroxy-16-methyl-9-oxoprost-13-en-1-oic acid methyl ester

Molecular formula: C₂₂H₃₈O₅

Molecular weight: 382.5

Structural formula:



Physicochemical properties:
Misoprostol is a synthetic prostaglandin E₁ analog. It is a colorless to yellow, viscous liquid with a musty odor.

pH: 6.2

Melting Point: None Observed

Aqueous Solubility at 25°C:

<u>Solvent</u>	<u>pH</u>	<u>g/mL</u>
Water	6.2	0.3
HCl, 0.01M	2.0	0.3
Acetate, 0.01M	4.5	0.3
Phosphate, 0.01M	7.0	0.3

Organic Solubility at Ambient Temperature:

<u>Solvent</u>	<u>g/mL</u>
Ethanol	>10
Corn Oil*	>1
Soy Bean Oil*	>1
Cottonseed Oil*	>1

*Containing 10% v/v absolute ethanol

CLINICAL TRIALS

Large meta-analyses of randomized clinical trials show that diclofenac is associated with an increased incidence of stroke, cardiovascular death, and death from any cause when compared with placebo. Data also suggest that diclofenac, particularly when used at a high dose (150 mg daily), may have a higher risk of thrombotic CV events than other NSAIDs.

Large population-based observational studies, meta-analyses and systematic reviews suggest that diclofenac use is associated with an increased incidence of cardiovascular thrombotic events, including myocardial infarction and ischemic stroke. Results of some studies suggest that the CV risk is related to the dose and duration of diclofenac exposure and is greater in patients with risk factors for CV disease.

In two multicentre, double-blind, controlled clinical trials of 12 weeks duration involving a total of 346 and 339 patients with rheumatoid arthritis respectively, patient global assessments of the

arthritic condition revealed no statistically significant differences between diclofenac sodium 50 mg plus misoprostol and a fixed-combination of diclofenac/placebo.

In two multicentre, double-blind, controlled trials of four weeks duration in 455 and 361 patients with osteoarthritis, patient global assessments of the arthritic condition revealed no overall differences between diclofenac sodium 50 mg plus misoprostol and diclofenac/placebo.

A multicentre, double-blind, controlled trial of 6 weeks duration involving a total of 572 patients (154 in the diclofenac group, 152 in the diclofenac sodium 50 mg plus misoprostol group, 175 in the diclofenac sodium 75 mg plus misoprostol group and 91 in the placebo group) showed that diclofenac sodium 50 mg plus misoprostol three times daily and diclofenac sodium 75 mg plus misoprostol twice daily were equivalent to diclofenac/placebo in relieving the signs and symptoms of osteoarthritis.

A multicentre, double-blind, controlled trial of 12 weeks duration involving a total of 380 patients (107 in the diclofenac group, 107 in the diclofenac sodium 50 mg plus misoprostol group, 111 in the diclofenac sodium 75 mg plus misoprostol group and 55 in the placebo group) showed that diclofenac sodium 50 mg plus misoprostol three times daily and diclofenac sodium 75 mg plus misoprostol twice daily were equivalent to diclofenac/placebo in relieving the signs and symptoms of rheumatoid arthritis.

Misoprostol has been compared to placebo in the prevention of clinically significant and serious gastrointestinal events associated with NSAID use. In a six-month, double-blind study of 8,843 patients (4,404 in the misoprostol group, 4,439 in the placebo group, mean age 68 years) with rheumatoid arthritis, misoprostol significantly reduced the incidence of serious complications, such as gastrointestinal bleeding and ulcer perforation, by 40-50%.

Diclofenac sodium plus misoprostol is associated with a low incidence of gastroduodenal lesions relative to diclofenac/placebo.

DETAILED PHARMACOLOGY

Diclofenac inhibits cyclo-oxygenase and thus reduces prostaglandin synthesis. This inhibition is believed to underlie the anti-inflammatory and analgesic effects of diclofenac. In addition, evidence suggests that increases in plasma beta-endorphin concentrations may mediate diclofenac-induced analgesia. The inhibition of prostaglandin synthesis is also believed to be involved in diclofenac-induced gastroduodenal mucosal injury. Increases in fecal blood loss and endoscopically verified mucosal damage were associated with diclofenac, as were increases in gastric transmural potential difference (i.e., an index of gastric irritation). Diclofenac has also demonstrated significant inhibition of platelet aggregation induced by adenosine diphosphate (ADP), adrenalin and collagen. However, its effects on spontaneous aggregation appear to vary with the disease state of the patient. Increases in blood urea have been associated with diclofenac in patients with osteoarthritis, but decreases in proteinuria were seen in patients with membranoproliferative disease or IgA glomerulonephritis.

Misoprostol enhances several of the factors implicated in maintaining gastroduodenal mucosal integrity. Basal-, histamine- and meal-stimulated acid secretion are reduced by misoprostol.

Mucosal blood flow which supplies nutrients and oxygen to the gastric mucosa has been shown to be increased by misoprostol. Misoprostol increases bicarbonate secretion from the proximal and distal segments of the duodenum in a dose-related manner. Misoprostol also increases basal mucus concentration and pentagastrin-stimulated mucus output and concentration.

Coadministration of misoprostol with ASA did not significantly affect the antiplatelet effects associated with ASA alone. Misoprostol reduced the decrement in renal function observed in patients with alcoholic cirrhosis and ascites who were receiving indomethacin.

Misoprostol 200 µg coadministered with diclofenac 50 mg TID to healthy volunteers had no clinically significant antiplatelet effects and would present no additional risk of bleeding diathesis apart from that due to diclofenac alone. The concurrent administration of diclofenac 50 mg BID and misoprostol 200 µg BID did not produce a significant treatment difference for whole bowel transit time, stool frequency or stool consistency from administration of either of the individual compounds. Misoprostol did not exacerbate renal impairment in rheumatoid arthritis patients with mild to moderate renal impairment who were receiving diclofenac for their arthritis.

TOXICOLOGY

Single-Dose Toxicity

Diclofenac Sodium

Species	Route	LD ₅₀ Range (mg/kg)
Mouse	p.o.	185-541
	i.v.	92-147
Rat	p.o.	55-240
	i.v.	97-161
Rabbit	p.o.	125-300
Dog	p.o.	>800
Monkey	p.o.	3200

The main clinical signs included convulsions, saltatory spasms, reduced activity, diarrhea and signs of acute systemic illness. The oral LD₅₀ for the dog was >800 mg/kg and for the monkey 3200 mg/kg. Dogs had transient anorexia, diarrhea and duodenal erosions. Monkeys had diarrhea, anorexia, emesis, salivation and rectal ulcers.

Misoprostol

Species	Route	LD ₅₀ Range (mg/kg)
Mouse	p.o.	27-138
	i.p.	70-160
Rat	p.o.	81-100
	i.p.	40-62
Dog	p.o.	9.0*

*single value

The main clinical signs were reduced activity and diarrhea in the mouse and rat. The main clinical signs in the dog were emesis, diarrhea, tremors and mydriasis.

Diclofenac Sodium/Misoprostol

(250:1; diclofenac sodium/misoprostol)

Species	Sex	Route	*LD ₅₀ Range (mg/kg)
Mouse	M	p.o.	110-190
	F	i.p.	140-240
Rat	M	p.o.	220-490
	F	i.p.	110-180

* Expressed as dosage of diclofenac sodium

Repeat-Dose Toxicity

Diclofenac Sodium

Studies in Rats with Diclofenac Sodium:

Studies of 4, 13, 15 and 26 weeks were done either in Wistar or Sprague-Dawley strains.

All rats in an oral 4-week study at doses of 0.5 to 16 mg/kg died within seven days. The main findings were necrosis (ulceration of the gastrointestinal mucosa, peritoneal adhesions, hypertrophy of mesenteric nodes and spleen, hemorrhages and hypoplasia of bone marrow). No deaths occurred in a diet admix study at doses up to 2 mg/kg, and there were no gastrointestinal ulcers. There were no adverse effects in one subcutaneous study at doses up to 6 mg/kg but in another subcutaneous study, one female at 10 mg/kg died and three had intestinal ulcers. Two females at the 6 mg/kg dose also had ulcers.

In the 15-week and 90-day studies with 4 and 6 mg/kg, respectively, there were significant decreases in hemoglobin, packed cell volume, total protein and alpha globulin. Significant increases occurred in reticulocytes and neutrophils. No gastrointestinal lesions were found in the 13-week study, but one 6 mg/kg rat in the 15-week study had fibrinopurulent peritonitis.

In a 6-month rat study at doses from 0.25 to 4 mg/kg, no effects occurred up to 1 mg/kg. At 2 mg/kg, females had an increase in neutrophils and hypertrophied mesenteric nodes. At 4 mg/kg, females had lowered hemoglobin, hematocrit and erythrocyte counts and an increase in leucocytes and neutrophils. Some also had intestinal ulcers with peritonitis and enlarged mesenteric nodes.

In a 98-week diet admix study in rats at doses of 0.25, 1.0 and 2.0 mg/kg, the high dose was terminated at 59 weeks because of deaths; 54% in males and 92% in females. Rats in the medium and high-dose groups had lowered hemoglobin, hematocrit and erythrocyte counts, and neutrophilic leucocytosis. Females at 1 and 2 mg/kg had lowered serum glucose and raised serum alkaline phosphatase activity. Livers of the four surviving high-dose females at week 59 weighed about 1.5 times more than control livers. Adrenal weight for medium and high-dose females was significantly increased.

Drug-related morphologic changes included splenomegaly, enlarged mesenteric nodes, ulceration of the small intestine, peritonitis (mainly in 1.0 and 2.0 mg/kg dose females), adrenocortical atrophy, prostatitis and plasma cell hyperplasia of mesenteric nodes.

Studies in Dogs with Diclofenac Sodium:

Studies of 16, 30 and 90 days were done by oral doses ranging from 0.5 to 10 mg/kg. Drug-related deaths or severe illness characterized by weakness and weight loss occurred at doses as low as 1.0 mg/kg. Findings were as follows: lowering of hemoglobin, hematocrit, erythrocyte count, total protein and albumin, elevation of leucocyte and reticulocyte counts and alpha and beta globulins, and marked hematopoiesis in the spleen with splenomegaly.

Changes were associated with erosions and perforated ulcers of the gastric and duodenal mucosa. No changes were reported at the low dose of 0.5 mg/kg in the 90-day study.

Studies in Primates with Diclofenac Sodium:

In a 3-month study in Rhesus monkeys at 5, 15 and 50 mg/kg and in a 6-month study at 5, 15 and 75 mg/kg, deaths occurred only at 75 mg/kg. Diarrhea/loose stools occurred at all doses. Hemoglobin, hematocrit and erythrocyte counts were decreased at all doses, while platelets, leucocytes and reticulocytes were increased. There were no gross or microscopic changes at doses up to 50 mg/kg. At 75 mg/kg the following were noted: alkaline phosphatase and blood urea nitrogen levels were elevated, total protein was decreased, liver weight was increased with signs of cellular vacuolation and hypertrophy, cellular vacuolation of kidneys with hyaline casts and debris in tubules and erosions/ulcers and hemorrhage in the gastrointestinal tract.

A one-year oral toxicity study was done in baboons at doses of 5, 15 and 50 mg/kg/day. Administration of the high dose was discontinued on day 19 and reinitiated on day 38 at a lower dose of 30 mg/kg. The medium dose was reduced from 15 mg/kg to 10 mg/kg on day 254. Five of 16 medium and 15 of 16 high-dose animals died or were killed in extremis. The main signs were emesis, shivering, lethargy, skin ulcers, bloody feces, facial edema and greatly reduced weight gain. At the medium and high doses, there was a dose-related decrease in hemoglobin, hematocrit value and erythrocyte count, and an increase in reticulocytes, neutrophilic leucocytes and platelets, with a left shift in the differential count. Serum globulins were increased. Shallow colonic ulcers were seen at the low dose. At the high dose, there were gastrointestinal ulcers,

some of which had perforated with resultant peritonitis. Gastrointestinal changes at the low and medium doses were less severe. Animals allowed to recover had no drug-related lesions of the stomach or intestines.

Misoprostol

Studies in Rats with Misoprostol:

Two 4, 5, 13 and 52 weeks toxicity studies were done in rats at daily oral dosages up to 9,000 µg/kg. There were no drug related deaths.

The clinical signs were diarrhea, salivation, vaginal dilation and discharge, decreased body weight gain and increased food consumption.

In the 52-week study, there were no abnormal clinical signs at 160 µg/kg and all signs at the higher doses were absent at the end of a 13-week reversal period.

Clinical laboratory changes included decreases in serum total protein and increases in serum iron. Study serum total protein decreased approximately 7 to 11% at 9,000µg/kg.

Hyperkeratosis of the aglandular part of the stomach and mucosal epithelial hyperplasia of the glandular part were the prominent gross and microscopic changes at all dosages. Hyperplasia of the superficial epithelial cells of the colon was observed at 9,000 µg/kg, but were absent at the end of the reversal period. The morphologic changes in the stomach were reflected in increased stomach weights and stomach to body weight ratios.

Electron microscopy of the stomach mucosa of some control and high dose (9,000µg/kg) animals showed the aglandular part of the stomach of treated animals had hyperkeratosis on the mucosal surface but the mucosal cells and keratin had normal structure. The corpus and antrum of the high dose rats had increased depth of gastric pits.

Studies in Dogs with Misoprostol:

Two 5, 13, and 52-week toxicity studies were conducted in beagle dogs at daily oral dosages ranging from 30 to 1,000µg/kg/day. The most prominent clinical signs were emesis, diarrhea, soft and/or mucoid stools and increased rectal temperatures.

The clinical observations were absent or decreased in severity at the end of the reversal periods (13 and 52 week studies).

One animal was killed *in extremis* at 300µg/kg during the first week of the study because it had stopped eating.

In the 52-week study, mean chloride concentrations were significantly increased, approximately 2, 4 and 5% in females at 30, 100 and 300µg/kg dosages, respectively. There were no abnormal clinical laboratory findings at the end of the reversal periods.

Radiographic examination of long bones was performed after 10 months in the 52-week toxicity and showed no significant differences between misoprostol-treated and control animals. There was no evidence of hyperostosis.

Reversible gastric mucosal epithelial hyperplasia was a consistent gross and microscopic change.

After a four-week recovery period in the 13-week study, a slight villous epithelial hyperplasia remained in the 480 µg/kg group. After a three-month recovery period in the 52-week study, there were no gross changes in the stomach and only one 300 µg/kg group male dog had hyperplasia of the pyloric epithelium.

Diclofenac Sodium/Misoprostol

Species	Period	No Significant Toxic Effect (mg/kg)*	Minimum Lethal Dose(mg/kg)*
Rat	4 weeks	>6	>6
	6 months	1	6
Dog	4 weeks	0.5	2
Monkeys	6 months	>6	50

*Expressed as dosage of diclofenac sodium (p.o. daily dose)

Studies in Rats with Diclofenac Sodium/Misoprostol:

In a 4-week oral study in rats, the dosages of misoprostol and diclofenac sodium, respectively, were 2, 8 and 24 µg/kg and 0.5, 2 and 6 mg/kg. Two other groups were given diclofenac sodium alone at 0.5 and 6 mg/kg. There were no deaths, clinical signs and gross and microscopic changes. Serum albumin concentrations were decreased in all treated females at all doses.

In a 6-month oral study in rats, the dosages of misoprostol and diclofenac sodium, respectively, were 4, 10, and 24 µg/kg and 1, 2.5, and 6 mg/kg. Three other groups were given diclofenac sodium alone at 1, 2.5 and 6 mg/kg. Gross and microscopic findings, seen primarily in the females of the high dosage groups, included jejunal ulceration, frequently associated with a granulation tissue reaction, jejunal dilatation and/or thickening, and diffuse peritonitis.

The splenic enlargement and extramedullary hematopoiesis of the spleen were considered secondary to ulceration. Jejunal ulcers were present in two females of the high combination group after the 4-week reversal period.

Hematological and serum chemical changes included anemia, thrombocytosis, neutrophilic leucocytosis, decreased serum protein and albumin, and increased alkaline phosphatase. Most of the changes were still present at the end of the reversal period. These changes were considered consistent with direct or indirect ulcerogenic effects of diclofenac on the gastrointestinal tract.

Studies in Dogs with Diclofenac Sodium/Misoprostol:

In a 4-week oral study in dogs, the dosages of misoprostol and diclofenac sodium, respectively, were 2, 4 and 8 µg/kg and 0.5, 1 and 2 mg/kg. Two other groups were given diclofenac sodium alone at 0.5 and 2 mg/kg. One dog died at the high diclofenac dose and three at the high

combined dose. These deaths were attributed to the effects of diclofenac.

Clinical signs in the two high dose groups included dark or bloody feces, loose stools, emesis, reduced activity and pale mucous membranes. Gross and microscopic findings included gastrointestinal ulcers, peritonitis, necrosis/edema of the renal crest, increased splenic hematopoiesis, and thymic and prostatic atrophy. Dogs in the two high dose groups were anemic and had neutrophilic leucocytosis, lowered plasma protein, albumin and calcium. In the two high dose groups mean splenic weight was increased and mean thymic weight decreased.

Diclofenac induced injury to the renal crest (hyperemia, edema and necrosis) in equal incidence whether given alone or in combination with misoprostol. There was a significant difference in severity of the lesions with necrosis occurring in 1/8 animals given the combination compared to 3/8 animals given diclofenac alone.

Studies in Monkeys with Diclofenac Sodium/Misoprostol:

In a 26-week oral study in *Cynomolgus* monkeys, the dosages of misoprostol and diclofenac sodium, respectively, were 24, 68, and 200 µg/kg and 6, 17, and 50 mg/kg. Two other groups were given diclofenac sodium alone at 6 and 50 mg/kg. Clinical signs of loose stools and salivation were mainly observed in combination groups and were considered to be related to the effect of misoprostol.

Significantly decreased body weight gains were observed in high-diclofenac males starting at week 3 and were not present at the end of the reversal period. Gross and microscopic changes including hyperemia, hemorrhages, and ulcerations of the gastrointestinal tract with various inflammatory reactions in other organs including peritonitis were seen in animals dying during the study and in animals from both high dose groups at the scheduled 26-week sacrifice. In a high-diclofenac female, acute myocarditis and pericarditis were associated with suppurative pyelonephritis. After 4 weeks reversal, cecal mucosal hemorrhage was still observed in two high combination males and one male each from the low and high-diclofenac groups.

Clinical laboratory changes were observed in animals of both high dose groups, but were generally more marked in animals treated with diclofenac alone. These changes included: anemia with increased reticulocytes and neutrophil count and decreased lymphocytes; thrombocytosis; decreased serum albumin and calcium; and increased serum globulin. With the exception of the reticulocyte counts in the high dose diclofenac group, the clinical laboratory changes were no longer observed or were subsiding at the end of the reversal period.

Genotoxicity

Diclofenac Sodium

No evidence of mutagenic potential was found in the following test systems: Ames Salmonella/microsome assay and yeast (*S. cerevisiae*) mutation, mouse lymphoma TK^{+/-} assay, nucleus anomaly in Chinese hamster bone marrow cells, chromosomal aberration and dominant lethal.

Misoprostol

The mutagenic/carcinogenic potential of misoprostol was evaluated in five *in vitro* assays: Ames

Salmonella/microsome assay; mouse lymphoma TK^{+/-} assay; sister chromatid exchange assay; yeast gene conversion assay; and the C₃H 10T1/2 cell transformation assay. Misoprostol was negative in all tests. Ames tests were also negative for misoprostol degradation products (SC-29636, SC-32759, SC-33188).

Diclofenac Sodium/Misoprostol

No evidence of mutagenicity was found in the following test systems: Ames Salmonella/microsome assay, CHO/HGPRT mutation assay, *in vitro* chromosome aberration assay in rat lymphocytes and in the mouse bone marrow micronucleus test.

Carcinogenicity

Carcinogenicity studies were conducted with misoprostol in rats and mice and with diclofenac sodium in rats. Neither diclofenac nor misoprostol is carcinogenic. Carcinogenicity studies have not been conducted with the combination of diclofenac sodium and misoprostol.

Reproductive and Developmental Toxicity

Reproductive Studies:

Diclofenac Sodium

Fertility (Segment I) and perinatal/postnatal (Segment III) studies were performed in the rat and teratology (Segment II) studies were performed in the mouse, rat and rabbit. There was no effect on fertility but maternal toxicity (intestinal ulcers and peritonitis) was produced at the high dose of 4 mg/kg. Postnatal growth and survival of pups in the Segment I study were equal between treated and control groups.

No teratogenic effects were evident in any of the Segment II studies but maternotoxicity and embryotoxicity occurred in some. Diclofenac has been shown to cross the placental barrier in mice and rats. Maternal deaths occurred at both doses of 2 and 4 mg/kg in the Segment III study. All dams that died or were killed *in extremis* had peritonitis, presumably associated with intestinal ulcers. Stillbirths and resorptions were higher in the two treated groups. Pups from dams receiving 4 mg/kg grew at a slower rate after the first week. Viability of offspring from surviving dams was not reduced.

Misoprostol

Fertility (Segment I) and perinatal/postnatal (Segment III) studies in rat and teratology (Segment II) studies in rat and rabbit were performed. There were drug-related clinical signs of salivation, soft feces, lethargy, and unkempt appearance at the higher doses of misoprostol. At a dosage of 100 µg/kg, no drug-related clinical signs occurred.

Although no drug-related deaths occurred, at doses of 1,600 µg/kg and above in rats and 300 µg/kg and above in rabbits decreases in body weights of male or female animals given misoprostol were observed.

In two rat fertility studies the number of implantations was decreased at 1,600 µg/kg and above. An increased number of resorptions occurred at 1,000 and 10,000 µg/kg in one study but were not reproduced in other studies. The increased number of resorptions and decreased number of implantations accounted for a decreased number of live fetuses or pups at 10,000 µg/kg, whereas

the decreased number of implantations accounted for a decreased number of fetuses at 1,600 µg/kg. Fetal and pup survival or growth were unaffected. Behavioral, sensory, and reproductive assessment of the F1 offspring revealed no adverse effects.

There was no evidence of embryotoxicity, fetotoxicity, or teratogenicity in two teratology rat studies at the maximum dosage of 10,000 µg/kg.

No evidence of fetotoxicity or teratogenicity was observed in two teratology rabbit studies at the maximum dosage of 1,000 µg/kg. However, there was an increased number of resorptions, evidence of possible embryotoxicity, in one of the two studies at 1,000 µg/kg.

In the perinatal/postnatal study, pup growth at 10,000 µg/kg was retarded as evidenced by the decreased weight gain during lactation. However, pup survival was unaffected.

Diclofenac Sodium/Misoprostol

One oral teratology (Segment II) study was done in rabbits. The dosages of misoprostol and diclofenac sodium, respectively, were 4, 12 and 40 µg/kg and 1, 3 and 10 mg/kg. At the high dose, there was lowered food intake, decreased weight gain, embryotoxicity and one possibly drug-related death.

At the lower doses, neither embryotoxicity nor maternotoxicity was noted. There was no evidence at any dose of fetotoxicity or teratogenicity.

REFERENCES

NSAIDs

1. Armstrong CP, Blower AL. Nonsteroidal anti-inflammatory drugs and life threatening complications of peptic ulceration. *Gut* 1987;28:527-532.
2. Champion MC. NSAID-associated mucosal damage and ulceration - Incidence, etiology, prophylaxis and treatment. *Can J Gastroenterol* July 1993;7(5):427-439.
3. Larkai EN, Smith JL, Lidsky MD, Graham DY. Gastroduodenal mucosa and dyspeptic symptoms in arthritic patients during chronic nonsteroidal anti-inflammatory drug use. *Am J Gastroenterol* 1987;82:1153-1158.
4. Somerville K, Faulkner G, Langman M. Non-steroidal anti-inflammatory drugs and bleeding peptic ulcer. *Lancet* 1986;1:462-464.

DICLOFENAC

1. Brogden RN, Heel RC, Pakes GE, Speight TM, Avery GS. Diclofenac Sodium: A review of its pharmacological properties and therapeutic use in rheumatic diseases and pain of varying origin. *Drugs* 1980;20:24-48.
2. Todd PA, Sorkin EM. Diclofenac sodium: A reappraisal of its pharmacodynamic and pharmacokinetic properties, and therapeutic efficacy. *Drugs* 1988;35:244-285.
3. Willis, JV, Kendall, MJ, Flinn, RM, Thornhill, DP, Welling PG. The pharmacokinetics of diclofenac sodium following intravenous and oral administration. *Eur J Clin Pharmacol* 1979;16:405-410.

MISOPROSTOL

1. Dajani EZ, Nissen CH. Gastrointestinal cytoprotective effects of misoprostol: clinical efficacy overview. *Dig Dis Sci* 1985;30:194S-200S.
2. Graham DY, Agrawal NM, Roth SH. Prevention of NSAID-induced gastric ulcer with misoprostol: multicentre, double-blind, placebo-controlled trial. *Lancet* 1988;3:1277-1280.
3. Karim A. Antiulcer prostaglandin misoprostol: single and multiple dose pharmacokinetic profile. *Prostaglandins* 1987;33(Suppl):40-50.
4. Steiner JA. Misoprostol clinical pharmacology: establishment of activity in man. *Dig Dis Sci* 1985;30:136S-141S.
5. Roth S, Agrawal N, Mahowald M, Montoya H, Robbins D, Miller S, Nutting E, Woods E, Crager M, Nissen C, Swabb E. Misoprostol heals gastroduodenal injury in patients with rheumatoid arthritis receiving aspirin. *Arch Int Med* 1989;149:775-779.

6. Silverstein FE, Graham DY, Senior JR, Davies HW, Struthers BJ, Bittman RM, Geis GS. Misoprostol reduces serious gastrointestinal complications in patients with rheumatoid arthritis receiving nonsteroidal anti-inflammatory drugs. *Annals of Internal Medicine*, August 15, 1995;123(4):241-249.

DICLOFENAC SODIUM/MISOPROSTOL

1. Geis GS, Stead H, Wallemark C, Nicholson PA. Prevalence of mucosal lesions in the stomach and duodenum due to chronic use of NSAID in patients with rheumatoid arthritis or osteoarthritis, an interim report on prevention by misoprostol of diclofenac associated lesions. *J Rheumatol* 1991;18:11-14.
2. Stead H, Geis GS. Treatment of NSAID-induced gastric and duodenal ulcers with misoprostol in patients with rheumatoid arthritis or osteoarthritis. *Brit J Rheum* 1992;31 (Suppl 2):177.
3. Geis GS, Erhardt LJ, Stead H. Prevention of diclofenac-induced gastroduodenal mucosal ulcers by misoprostol: a one year study. *Brit J Rheum* 1992;31 (Suppl 2):180.

GD-diclofenac/misoprostol (ARTHROTEC)

1. Verdickt W, Moran C, Hantzschel H, Fraga AM, Stead H, Geis GS. A double-blind comparison of the gastroduodenal safety and efficacy of diclofenac and a fixed dose combination of diclofenac and misoprostol in the treatment of rheumatoid arthritis. *Scand J Rheum* 1992;21:85-91.
2. Bolten W, De Melo Gomes JA, Stead H, Geis GS. The gastroduodenal safety and efficacy of the fixed combination of diclofenac and misoprostol in the treatment of osteoarthritis. *Brit J Rheum* 1992;31:753-758.
3. Hannequin JR. Efficacy of Arthrotec in the treatment of rheumatoid arthritis. *Scand J Rheumatol* 1992;Suppl 96:7-14.
4. Doherty M. The efficacy of Arthrotec in the treatment of osteoarthritis. *Scand J Rheumatol* 1992; Suppl 96:15-21.
5. De Melo Gomes JA. The safety of Arthrotec in patients with rheumatoid arthritis or osteoarthritis: An assessment of the upper gastrointestinal tract by endoscopy. *Scand J Rheumatol* 1992; Suppl 96:23-31.
6. Geis GS. Overall safety of Arthrotec. *Scand J Rheumatol* 1992; Suppl 96:33-36.
7. De Melo Gomes JA, Roth SH, Zech J, Bruyn GAW, Woods EM, Geis GS. Double-blind comparison of efficacy and gastroduodenal safety of diclofenac/misoprostol, piroxicam, and naproxen in the treatment of osteoarthritis. *Annals of the Rheumatic Diseases* 1993;52:881-885.
8. Karim A, Smith M. Biopharmaceutical profile of diclofenac-misoprostol combination tablet, Arthrotec. *Scand J Rheumatol* 1992;Suppl 96:37-48.

9. Kiff PS, Stead H, Morant SV, Shield MJ. Arthrotec, diclofenac and ibuprofen in general practice. *European J Rheumatol* 1994;14(Issue 3):31-38.
10. De Queriroz MV, Beaulieu A, Kruger MD, Woods E, Stead H, Geis GS. Double-blind comparison of the efficacy of diclofenac/misoprostol and diclofenac in the treatment of rheumatoid arthritis. *Eur J Rheumatol Inflamm* 1994;14(Issue 2):5-13.

PART III: CONSUMER INFORMATION

PrGD-diclofenac/misoprostol 50
PrGD-diclofenac/misoprostol 75

diclofenac sodium and misoprostol enteric-coated tablets

This leaflet is part III of a three-part "Product Monograph" published when GD-diclofenac/misoprostol was approved for sale in Canada and is designed specifically for Consumers.

Read this information each time you refill your prescription in case new information has been added.

This leaflet is a summary and will not tell you everything about GD-diclofenac/misoprostol. See your health care provider and pharmacist regularly and ask them questions about your health and any medications you take.

ABOUT THIS MEDICATION

What the medication is used for:

Your health care provider has prescribed GD-diclofenac/misoprostol for you for the following:

- short-term and long-term use in the relief of the signs and symptoms of rheumatoid arthritis;
- short-term and long-term use in the relief of the signs and symptoms of osteoarthritis

What it does:

GD-diclofenac/misoprostol contains two different medicines, a nonsteroidal anti-inflammatory drug (NSAID) called diclofenac and a drug that helps to protect the lining of your stomach called misoprostol (because NSAIDs can cause damage to your stomach).

GD-diclofenac/misoprostol (diclofenac plus misoprostol), as a nonsteroidal anti-inflammatory drug (NSAID), can reduce the chemicals produced by your body which cause pain and swelling. It helps to relieve joint pain, swelling, and stiffness by reducing the production of certain substances (prostaglandins) and by helping to control inflammation. GD-diclofenac/misoprostol, as a nonsteroidal anti-inflammatory drug (NSAID) does NOT cure your illness or prevent it from getting worse but it promotes suppression of the inflammation and the tissue damaging effects resulting from this inflammation. GD-diclofenac/misoprostol can only relieve pain and reduce swelling as long as you continue to take it.

How do NSAIDs cause stomach damage? Natural prostaglandins play an important role in protecting the stomach by working to keep a thick mucus layer on the inside surface of the stomach. If the lining of the stomach is not protected by a thick layer of mucus, it may be burned by natural stomach acids. NSAIDs lower natural prostaglandins both in the joints and in the stomach. This is good for the joints because it controls pain, swelling and stiffness. Unfortunately, lowering prostaglandins in the stomach can lead to burning stomach pain and the development of tiny holes in the lining of your stomach called "ulcers".

Oddly enough, some NSAID patients who do develop ulcers never feel any stomach pain. On the other hand, some patients who do feel stomach pain have nothing wrong with them. For this reason people must be very careful when taking NSAIDs.

How does misoprostol protect the stomach? Misoprostol is a synthetic form of a special kind of prostaglandin that is found in the stomach. Misoprostol replaces the prostaglandins that are lost when taking the NSAID medicine. It protects the thick mucus layer and reduces the acid in your stomach. This can help to protect your stomach from the NSAID.

When it should not be used:

SPECIAL NOTE FOR WOMEN OF CHILDBEARING AGE

Do not take GD-diclofenac/misoprostol if you are pregnant or think you are pregnant. Do not start GD-diclofenac/misoprostol until you have been tested to confirm that you are not pregnant. You should avoid becoming pregnant while you are taking GD-diclofenac/misoprostol and for at least one month (or through one menstrual cycle) after you stop taking it. This means using an effective form of birth control which you should discuss with your doctor. Stop taking GD-diclofenac/misoprostol, and contact your doctor immediately if you do become pregnant during GD-diclofenac/misoprostol therapy.

Misoprostol may cause uterine contractions (contractions of the uterus), premature birth, birth defects and abortion or may otherwise harm the unborn developing baby. Misoprostol has been reported to cause the uterus to tear when given after the eighth week of pregnancy. Tearing of the uterus can result in severe bleeding, hysterectomy, and/or maternal or fetal death. Therefore, if you are pregnant, you must not take this drug.

Abortions caused by misoprostol are likely to be incomplete. An incomplete abortion may result in very serious medical complications, resulting in hospitalization, surgery and possibly infertility, and may result in maternal death.

You should not take GD-diclofenac/misoprostol if you are nursing. The body changes misoprostol to the active form of the drug, misoprostol acid, which could get into the breast milk and cause significant diarrhea in the infant.

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

DO NOT TAKE GD-diclofenac/misoprostol if you have any of the following medical conditions:

- Heart bypass surgery (planning to have or recently had)
- Severe, uncontrolled heart failure
- Bleeding in the brain or other bleeding disorders
- Current pregnancy or pregnancy cannot be excluded
- Currently breastfeeding (or planning to breastfeed)
- Allergy to diclofenac sodium, misoprostol, ASA (Acetylsalicylic Acid) or other NSAIDs (Nonsteroidal Anti-Inflammatory Drugs)
- Ulcer (active)
- Bleeding from the stomach or gut (active)
- Inflammatory bowel disease (Crohn's Disease or Ulcerative Colitis)
- Liver disease (active or severe)
- Kidney disease (severe or worsening)
- High potassium in the blood
- Allergy to any other ingredients in **GD-diclofenac/misoprostol** listed in the "nonmedicinal ingredients" section below

Patients who took a drug in the same class as GD-diclofenac/misoprostol after a type of heart surgery [coronary artery bypass grafting (CABG)] were more likely to have heart attacks, strokes, blood clots in the leg(s) or lung(s), and infections or other complications than those who did NOT take that drug.

GD-diclofenac/misoprostol should NOT be used in patients under 18 years of age since safety and effectiveness have NOT been established.

GD-diclofenac/misoprostol should NOT be used with other NSAIDs.

What the medicinal ingredients are:

Diclofenac sodium and misoprostol

What the important nonmedicinal ingredients are:

Lactose, Hydrogenated Castor Oil, Microcrystalline Cellulose, Colloidal Silicon Dioxide, Corn Starch, Crospovidone, Hypromellose, Magnesium Stearate, Methacrylic Acid Copolymer, Povidone K-30, Sodium Hydroxide, Talc, Triethyl Citrate.

What dosage forms it comes in:

Enteric coated Tablets:

GD-diclofenac/misoprostol 50: 50 mg diclofenac sodium/200 µg misoprostol,

GD-diclofenac/misoprostol 75: 75 mg diclofenac sodium/200 µg misoprostol

WARNINGS AND PRECAUTIONS

If you have, or previously had, any of the following medical conditions, see your health care provider to discuss treatment options other than GD-diclofenac/misoprostol:

- Heart Attack or Angina
- Stroke or Mini-stroke
- Loss of Vision
- Congestive Heart Failure
- Current Pregnancy
- High blood pressure
- Diabetes
- High levels of fats in your blood
- Smoking

It is important to take the lowest dose of GD-diclofenac/misoprostol that relieves your pain and/or swelling and for the shortest time possible in order to keep your risk of side effects on the heart and blood vessels as small as possible.

Use of NSAIDs, such as GD-diclofenac/misoprostol may cause stomach and bowel problems (such as ulceration, perforation, obstruction, and bleeding).

Use of GD-diclofenac/misoprostol can result in increased blood pressure and / or worsening of congestive heart failure.

BEFORE you use GD-diclofenac/misoprostol talk to your doctor or pharmacist if you have any of the following:

- Disease of the heart or blood vessels (also called cardiovascular disease, including uncontrolled high blood pressure, congestive heart failure, established ischemic heart disease, or peripheral arterial disease), as treatment with GD-diclofenac/misoprostol in these cases is not recommended.
- Risk factors for cardiovascular disease (see above) such as high blood pressure, abnormally high levels of fat (cholesterol, triglycerides) in your blood, diabetes, or if you smoke.
- Diabetes mellitus or on a low sugar diet
- Atherosclerosis
- Poor circulation to your extremities
- Smoker or ex-smoker
- Kidney disease or urine problems
- Previous ulcer or bleeding from the stomach or gut
- Previous bleeding in the brain
- Bleeding problems
- Family history of allergy to NSAIDs, such as acetylsalicylic acid (ASA), celecoxib, diclofenac, diflunisal, etodolac, fenoprofen, fluribiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, mefenamic acid, meloxicam, nabumetone, naproxen, oxaprozin, piroxicam,

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

rofecoxib, sulindac, tenoxicam, tiaprofenic acid, tolmetin, or valdecoxib (NOT a complete list)

- Family history of asthma, nasal polyps, long-term swelling of the sinus (chronic sinusitis) or hives.
- History of a severe drug hypersensitivity reaction including skin problems
- you are on any special diet, such as a low-sodium diet.
- you drink alcohol;

GD-diclofenac/misoprostol may cause sensitivity to sunlight

Fertility may be decreased. The use of

GD-diclofenac/misoprostol is not recommended in women who have difficulty conceiving.

WHILE taking GD-diclofenac/misoprostol:

- tell any other doctor, dentist, pharmacist or other health care professional that you consult or see, that you are taking this medication, especially if you are planning to have heart surgery;
- do NOT drink alcoholic beverages while taking this medication because you would be more likely to develop stomach problems;
- some NSAIDs may cause drowsiness or fatigue in some people taking them; be cautious about driving or participating in activities that require alertness if you are drowsy, dizzy or lightheaded after taking this medication;
- your regular medical checkups are essential.
- do not give GD-diclofenac/misoprostol to anyone else. It has been prescribed for your specific condition. GD-diclofenac/misoprostol may not be the correct treatment for another person, and would be dangerous if the other person were pregnant.
- fertility may be decreased. The use of GD-diclofenac/misoprostol is not recommended in women trying to get pregnant. In women who have difficulty conceiving, stopping GD-diclofenac/misoprostol should be considered.
- if you have cardiovascular disease or risks for cardiovascular disease, your doctor will periodically re-evaluate whether you should continue treatment with GD-diclofenac/misoprostol.

If, at any time while taking GD-diclofenac/misoprostol you experience any signs or symptoms of problems with your heart or blood vessels such as chest pain, shortness of breath, weakness, or slurring of speech, contact your doctor immediately.

INTERACTIONS WITH THIS MEDICATION

Talk to your health care provider and pharmacist if you are taking any other medication (prescription or non-prescription) such as any of the following (NOT a complete list):

- Acetylsalicylic Acid (ASA) or other NSAIDs (e.g. ASA, celecoxib, diclofenac, ibuprofen, indomethacin, ketorolac, meloxicam, naproxen)
- Antacids

- Antidepressants
 - Selective Serotonin Reuptake Inhibitors (SSRIs) (e.g. citalopram, fluoxetine, paroxetine, sertraline)
- Blood pressure medications
 - Diuretics (e.g. furosemide, hydrochlorothiazide)
 - ACE (Angiotensin converting enzyme) inhibitors (e.g. enalapril, lisinopril, perindopril, ramipril)
 - ARBs (angiotensin II receptor blockers) (e.g. candesartan, irbesartan, losartan, valsartan)
 - Beta-blockers
- Blood thinners (e.g. warfarin, ASA, clopidogrel)
- Corticosteroids (including glucocorticoids) (e.g. prednisone)
- Cyclosporin
- Digoxin
- Lithium
- Methotrexate
- Oral contraceptives
- Oral hypoglycemics (diabetes medications)
- Phenytoin (a medicine used to treat seizures)
- Tacrolimus
- Sulfapyrazone (a medicine used to treat gout)
- Voriconazole (a medicine used to treat fungal infections)

Do not take ASA (acetylsalicylic acid), ASA-containing compounds, ibuprofen or other drugs used to relieve symptoms of arthritis while taking GD-diclofenac/misoprostol unless directed to do so by your physician.

Your health care provider may prescribe low dose ASA (acetylsalicylic acid) as a blood thinner to reduce your risk of having a heart attack or stroke while you are taking GD-diclofenac/misoprostol. Take only the amount of ASA prescribed by your healthcare provider. You are more likely to upset or damage your stomach if you take both GD-diclofenac/misoprostol and ASA than if you took GD-diclofenac/misoprostol alone.

Using GD-diclofenac/misoprostol with a blood-thinner increases the risk of bleeding. This can occur in the stomach or anywhere.

Using GD-diclofenac/misoprostol with blood pressure drugs may increase your risk of kidney failure. This is more likely if you are elderly or dehydrated.

Do not take antacids that contain magnesium (because they can cause diarrhea) while you are taking GD-diclofenac/misoprostol. Ask your pharmacist to help you select a suitable brand.

Stomach problems may be more likely to occur if you drink alcoholic beverages. Therefore, do not drink alcoholic beverages while taking this medication.

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

PROPER USE OF THIS MEDICATION

Usual Dose for Patients 18 Years of Age and Older:

Medical Condition	Recommended daily dose	Maximum Dose (per day)*
Rheumatoid arthritis	50 mg twice a day	100mg
Osteoarthritis	50 mg twice a day	100mg

GD-diclofenac/misoprostol should be swallowed whole.

To lessen stomach upset, take GD-diclofenac/misoprostol immediately after a meal or with food or milk. Also, you should remain standing or sitting upright (i.e. do not lie down) for about 15-30 minutes after taking the medicine. This helps to prevent irritation that may lead to trouble swallowing.

Take GD-diclofenac/misoprostol only as directed by your health care provider. **Do NOT take more of it, do NOT take it more often and do NOT take it for a longer period of time than your healthcare provider ordered. If possible, you should take the lowest dose of this medication for the shortest time period.** Taking too much GD-diclofenac/misoprostol may increase the chance of unwanted and sometimes dangerous side effects, especially if you are an elderly patient, have other disease or take other medications.

If you will be using GD-diclofenac/misoprostol for more than 7 days, see your healthcare provider regularly to discuss whether this medicine is working for you and if it is causing you any unwanted effects. Be sure to take GD-diclofenac/misoprostol regularly as prescribed. In some types of arthritis, up to two weeks may pass before you feel the full effects of this medicine. During treatment, your doctor may decide to adjust the dosage according to your response to the medication.

This medication has been prescribed specifically for you. Do NOT give it to anyone else. It may harm them, even if their symptoms seem to be similar to yours.

GD-diclofenac/misoprostol is NOT recommended for use in patients under 18 years of age since safety and effectiveness have NOT been established.

Missed Dose:

If you miss a dose of GD-diclofenac/misoprostol, take the next dose at the regular time. It is important that GD-diclofenac/misoprostol be taken as prescribed. Try to remember to take GD-diclofenac/misoprostol at the appropriate time. Having a regular routine associated with taking your medicine will help.

Do not take a double dose.

Overdose:

Symptoms of overdose may include stomach pain, confusion, drowsiness, low muscle tone, shaking hands that you cannot control, seizures, shortness of breath, diarrhea, fever, rapid or pounding heartbeat, slow heartbeat, dizziness or fainting

Seek immediate medical attention if you think that you or anyone else may have taken too much GD-diclofenac/misoprostol. Do this even if there are no signs of discomfort or poisoning.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

GD-diclofenac/misoprostol may cause some undesirable reactions especially when used for a long time or in large doses. When these side effects occur, you may require medical attention. Report all symptoms or side effects to your health care provider.

GD-diclofenac/misoprostol may cause you to become drowsy or tired. Be careful about driving or participating in activities that require you to be alert. If you become drowsy, dizzy or light-headed after GD-diclofenac/misoprostol, do NOT drive or operate machinery.

GD-diclofenac/misoprostol may cause you to become more sensitive to sunlight. Any exposure to sunlight or sunlamps may cause sunburn, skin blisters, skin rash, redness, itching or discoloration, or vision changes. If you have reaction from the sun, check with your health care provider.

Check with your healthcare provider IMMEDIATELY if you develop chills, fever, muscle aches or pains, or other flu-like symptoms, especially if they occur before or together with a skin rash. These symptoms may be the first signs of a SERIOUS ALLERGIC REACTION to this medication.

If you are not getting any relief of your arthritis or if any problems develop, check with your doctor.

Elderly, frail, or debilitated patients often seem to experience more frequent or more severe side effects.

Stomach upset is one of the common problems with NSAIDs. If stomach upset (indigestion, nausea, vomiting, stomach pain or diarrhea) occurs and continues, contact your doctor.

Because misoprostol increases mucus production some patients experience diarrhea. Keep taking your GD-diclofenac/misoprostol. It is just a sign that the drug is working. Usually the diarrhea goes away in two to three days. If it is not gone after a week, check with your doctor.

While your body gets used to misoprostol you may feel a crampy pain in your stomach. Like the diarrhea it usually goes away in a few days. If it doesn't, check with your doctor.

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

090177e19572def4\Final\Final On: 05-Nov-2020 22:33 (GMT)

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom	STOP taking GD-diclofenac/misoprostol and get emergency medical attention IMMEDIATELY	STOP taking GD-diclofenac/misoprostol and talk your doctor or pharmacist
Bloody or black tarry stools, vomiting blood	✓	
Shortness of breath, wheezing, any trouble in breathing or tightness in the chest	✓	
Skin rash, hives or swelling, itching, bruising	✓	
Blurred vision or any visual disturbance	✓	
Any change in the amount or colour of your urine (red or brown)	✓	
Nausea, fatigue, lethargy, diarrhea, pruritus, yellow discoloration of the skin or eyes with or without itchy skin, right upper quadrant tenderness, and «flu-like» symptoms	✓	
Change in heart rate or rhythm, change in blood pressure, heart failure, blood clot	✓	
Infections: Sepsis (infection of the whole body) Lung infection	✓	
Seizure, shaking, loss of consciousness, strange dreams, change in mood or thoughts	✓	
Benign mass in the intestine, trouble with swallowing	✓	
Serious skin reactions: skin rash, blisters or breakdown of your skin, fever, swollen glands		✓

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom	STOP taking GD-diclofenac/misoprostol and get emergency medical attention IMMEDIATELY	STOP taking GD-diclofenac/misoprostol and talk your doctor or pharmacist
Any pain or difficulty experienced while urinating		✓
Swelling of the feet or lower legs; weight gain		✓
Vomiting or persistent indigestion, nausea, stomach pain or diarrhea		✓
Malaise, fatigue, loss of appetite		✓
Headaches, stiff neck		✓
Dizziness, light headedness		✓
Hearing problems		✓
Agranulocytosis: fever, chills. Sore throat or mouth sores. Flu-like symptoms. Weakness.		✓
Hemolytic Anemia: Fatigue and short of breath. Yellow skin or eyes. Dark urine. Abdominal pain, nausea, vomiting, loss of appetite.		✓
Increased sensitivity or nerve pain		✓

This is not a complete list of side effects. For any unexpected effects while taking GD-diclofenac/misoprostol, contact your doctor or pharmacist.

HOW TO STORE IT

Store at 15-25°C and protect from heat and humidity.
Do NOT keep outdated medicine or medicine no longer needed. Any outdated or unused medicine should be returned to your pharmacist.

Keep out of the reach of children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to:
Canada Vigilance Program
Health Canada
Postal Locator 1908C
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals may be obtained by contacting the sponsor, GenMed, a division of Pfizer Canada ULC, at:
1-800-463-6001

This leaflet was prepared by
GenMed, a division of Pfizer Canada ULC
17,300 Trans-Canada Highway
Kirkland, Quebec H9J 2M5

Last revised: November 2, 2020